



*The Alberta Lake Management Society  
Volunteer Lake Monitoring Program*

## Moose Lake Report

# 2021

Updated May 6, 2022

Lakewatch is made possible  
with support from:





# ALBERTA LAKE MANAGEMENT SOCIETY'S LAKEWATCH PROGRAM

LakeWatch has several important objectives, one of which is to collect and interpret water quality data from Alberta's Lakes. Equally important is educating lake users about aquatic environments, encouraging public involvement in lake management, and facilitating cooperation and partnerships between government, industry, the scientific community and lake users. LakeWatch reports are designed to summarize basic lake data in understandable terms for the widest audience, and are not meant to be a complete synopsis of information about specific lakes. Additional information is available for many lakes that have been included in LakeWatch, and readers requiring more information are encouraged to seek those sources.

ALMS would like to thank all who express interest in Alberta's aquatic environments, and particularly those who have participated in the LakeWatch program. These leaders in stewardship give us hope that our water resources will not be the limiting factor in the health of our environment.

If you require data from this report, please contact ALMS for the raw data files.



## ACKNOWLEDGEMENTS

The LakeWatch program is made possible through the dedication of its volunteers. A special thanks to Cody Fedun and Kellie Nichiporik for their commitment to collecting data at Moose Lake. We would also like to thank Keri Malanchuk and Brittany Onysyk, who were summer technicians in 2021. Executive Director Bradley Peter and Program Manager Caleb Sinn were instrumental in planning and organizing the field program. This report was prepared by Caleb Sinn and Bradley Peter.

BEFORE READING THIS REPORT, CHECK  
OUT [A BRIEF INTRODUCTION TO  
LIMNOLOGY](#) AT [ALMS.CA/REPORTS](#)

## MOOSE LAKE

Moose Lake is located 240 km northeast of Edmonton and 3.5 km west of the Town of Bonnyville. Moose Lake has over 64 km of irregular shoreline within a 40 km<sup>2</sup> lake surface area. The lake is comprised of four main bays with a maximum depth of 19 m and a mean depth of 5.6 m. A sounding (whole lake depth measurement) was last conducted in 1962.

The lake was once known by its French name Lac d'Original, which was inspired by the abundance of moose in the area.<sup>1</sup> In 1789, Angus Shaw established a trading post for the North West Company on the northwest shore of Moose Lake, one of the earliest European settlements known to Alberta. Later, in the early 1900's, French Canadian settlers began arriving in the area. In 1928, the railway was extended from St. Paul to Bonnyville.<sup>1</sup>



Moose Lake—photo by Elashia Young 2017

Moose Lake's abundance of natural resources was in high demand to supply a rapidly expanding population. Mink farming, agriculture, and three commercial fish-packing plants were in operation by 1936.<sup>1</sup> Walleye, northern pike, and yellow perch are the most popular sport fish; however, the lake also contains cisco, lake whitefish, burbot, suckers, and forage fish. Moose Lake is still heavily used, particularly on summer weekends. Shoreline development is intense and includes cottage subdivisions, campgrounds, and summer villages. Aquatic reeds fringe the shoreline, which is predominantly sheltered. Dominant emergent plants include bulrush (*Scirpus validus*) and cattail (*Typha latifolia*). Common submergent plants are pondweeds (*Potamogeton* spp.) and northern watermilfoil (*Myriophyllum sibiricum*). Moose Lake also provides excellent habitat to a variety of waterfowl, although residents are concerned that the current high population level of cormorants (*Phalacrocorax auritus*) in the region, are contributing to poor water quality conditions at Moose Lake.

The watershed area for Moose Lake is 808.01 km<sup>2</sup> and the lake area is 40.53 km<sup>2</sup>. The lake to watershed ratio of Moose Lake is 1:20. A map of the Moose Lake watershed area can be found at <http://alms.ca/wp-content/uploads/2016/12/Moose.pdf>. Moreover, multi-basin monitoring of Moose Lake was conducted in 2016 and 2017, the results of which can be found at [www.alms.ca](http://www.alms.ca). A phosphorus budget for the lake was completed by Associated Environmental in 2021.<sup>2</sup>

<sup>1</sup> Mitchell, P. and E. Prepas. 1990. Atlas of Alberta Lakes, University of Alberta Press. Retrieved from <http://sunsite.ualberta.ca/projects/alberta-lakes/>

<sup>2</sup> [https://laraonline.ca/wp-content/uploads/2021/12/Final\\_rpt\\_AE\\_MooseLakePbudget\\_Sep-2021.pdf](https://laraonline.ca/wp-content/uploads/2021/12/Final_rpt_AE_MooseLakePbudget_Sep-2021.pdf). Accessed May 2022.



## WATER CHEMISTRY

*ALMS measures a suite of water chemistry parameters. Phosphorus, nitrogen, and chlorophyll-*a* are important because they are indicators of eutrophication, or excess nutrients, which can lead to harmful algal/cyanobacteria blooms. One direct measure of harmful cyanobacteria blooms are Microcystins, a common group of toxins produced by cyanobacteria. See Table 2 for a complete list of parameters.*

The average total phosphorus (TP) concentration for Moose Lake was 54 µg/L (Table 2), falling into the eutrophic, or high productivity trophic classification. This value is in the mid-range of historical averages. TP was lowest early in the season at 31 µg/L during the June 25<sup>th</sup> sampling event, and peaked later in the season at 73 µg/L during the August 13<sup>th</sup> sampling event (Figure 1).

Average chlorophyll-*a* concentration in 2021 was 48.1 µg/L (Table 2), falling into the hypereutrophic, or very highly productive trophic classification. Chlorophyll-*a* was lowest during the June 25<sup>th</sup> sampling event at 23.0 µg/L, and increased later in the season and peaked at 63.7 µg/L on August 13<sup>th</sup> (Figure 1).

The average TKN concentration was 2.1 mg/L (Table 2) and varied from 1.8 mg/L to 2.3 mg/L, having the highest levels during the July and August sampling events (Figure 1). TKN was significantly positively correlated with chlorophyll-*a* ( $r = 0.97$ ,  $p = 0.03$ ).




Figure 1. Total Phosphorus (TP), Total Kjeldahl Nitrogen (TKN), and Chlorophyll-*a* concentrations measured four times over the course of the summer at Moose Lake.

Average pH was measured as 8.79 in 2021, buffered by high alkalinity (312 mg/L  $\text{CaCO}_3$ ) and bicarbonate (332 mg/L  $\text{HCO}_3^-$ ). Sulphate, bicarbonate and sodium were the dominant ions found. Together, with the other major ions, they contributed to a moderate conductivity of 925  $\mu\text{S}/\text{cm}$  (Figure 2, top; Table 2). Moose Lake is in the moderate to high range of ion levels compared to other LakeWatch lakes sampled in 2021 (Figure 2, bottom).




Figure 2. Average levels of cations (sodium =  $\text{Na}^{1+}$ , magnesium =  $\text{Mg}^{2+}$ , potassium =  $\text{K}^{1+}$ , calcium =  $\text{Ca}^{2+}$ ) and anions (chloride =  $\text{Cl}^{1-}$ , sulphate =  $\text{SO}_4^{2-}$ , bicarbonate =  $\text{HCO}_3^{1-}$ , carbonate =  $\text{CO}_3^{2-}$ ) from four measurements over the course of the summer at Moose Lake. Top) bars indicate range of values measured, and bottom) Schoeller diagram of average ion levels at Moose Lake (blue line) compared to 25 lake basins (gray lines) sampled through the LakeWatch program in 2021 (note log<sub>10</sub> scale on y-axis of bottom figure).

## METALS

*Metals will naturally be present in aquatic environments due to in-lake processes or the erosion of rocks, or introduced to the environment from human activities such as urban, agricultural, or industrial developments. Many metals have a unique guideline as they may become toxic at higher concentrations. Where current metal data are not available, historical concentrations for 27 metals have been provided (Table 3).*

Metals were measured at Moose Lake in 2021, and no metal exceeded the CCME guidelines for the protection of aquatic life, chronic (Table 3).

## WATER CLARITY AND EUPHOTIC DEPTH

*Water clarity is influenced by suspended materials, both living and dead, as well as dissolved colored compounds in the water column. During the melting of snow and ice in spring, lake water can become turbid (cloudy) from silt transported into the lake. Lake water usually clears in late spring, but then becomes more turbid with increased algal growth as the summer progresses. The easiest and most widely used measure of lake water clarity is the Secchi depth. Two times the Secchi depth equals the euphotic depth – the depth to which there is enough light for photosynthesis.*

The average euphotic depth of Moose Lake in 2021 was 2.96 m, corresponding to an average Secchi depth of 1.48 m (Table 2). The euphotic depth was very deep during the June 25<sup>th</sup> sampling event at 6.5 m depth, but was substantially shallower during all subsequent sampling events, remaining consistent at about 0.90 m (Figure 3).




Figure 3. Euphotic depth values measured four times over the course of the summer at Moose Lake in 2021.

## WATER TEMPERATURE AND DISSOLVED OXYGEN

*Water temperature and dissolved oxygen (DO) profiles in the water column can provide information on water quality and fish habitat. The depth of the thermocline is important in determining the depth to which dissolved oxygen from the surface can be mixed. Please refer to the end of this report for descriptions of technical terms.*

Surface temperatures of Moose Lake varied throughout the summer, with the July 14<sup>th</sup> sampling date having the warmest temperatures at 22.2°C (Figure 4a). The lake was weakly stratified during the June and July sampling events, and was mixed during the August and September sampling events.

Moose Lake was well oxygenated in the surface waters on all sampling dates, measuring above the CCME guidelines of 6.5 mg/L dissolved oxygen (Figure 4b). Oxygen levels decreased appreciably at depths corresponding to the thermocline on June 25<sup>th</sup> and July 14<sup>th</sup>, about 8 m or 7.5 m, respectively. Below the thermocline depth, the oxygen levels proceeded to anoxia (<1.0 mg/L oxygen) near and at lake bottom. Despite a lack of thermal stratification on August 13<sup>th</sup>, oxygen level decreased appreciably below 5 m depth on that sampling date.




Figure 4. a) Temperature (°C) and b) dissolved oxygen (mg/L) profiles for Moose Lake measured four times over the course of the summer of 2021.



## MICROCYSTIN

*Microcystins are toxins produced by cyanobacteria (blue-green algae) which, when ingested, can cause severe liver damage. Microcystins are produced by many species of cyanobacteria which are common to Alberta's Lakes, and are thought to be one of the most common cyanobacteria toxins. In Alberta, recreational guidelines for microcystin are set at 10 µg/L. Blue-green algae advisories are managed by Alberta Health Services. Recreating in algal blooms, even if microcystin concentrations are not above guidelines, is not recommended.*

Microcystin levels in Moose Lake fell below the recreational guideline of 10 µg/L during every sampling event in 2021 (Table 1), but the seasonal average is on the higher end of historical averages (Table 2). Even though low levels of microcystin were detected in composite samples, caution should always be observed when recreating around cyanobacteria.

Table 1. Microcystin concentrations measured four times at Moose Lake in 2021.

Date	Microcystin Concentration (µg/L)
25-Jun-21	0.66
14-Jul-21	6.37
13-Aug-21	6.07
10-Sep-21	1.46
Average	3.64

## INVASIVE SPECIES MONITORING

*Dreissenid mussels pose a significant concern for Alberta because they impair the function of water conveyance infrastructure and adversely impact the aquatic environment. These invasive mussels can change lake conditions which can then lead to toxic cyanobacteria blooms, decrease the amount of nutrients needed for fish and other native species, and cause millions of dollars in annual costs for repair and maintenance of water-operated infrastructure and facilities. Spiny water flea pose a concern for Alberta because they alter the abundance and diversity of native zooplankton, as they are aggressive zooplankton predators. Through over-predation, they will impact higher trophic levels such as fish. They also disrupt fishing equipment by attaching in large numbers to fishing lines.*

Monitoring involved sampling with a 63 µm plankton net at three sample sites to look for juvenile mussel veligers and spiny water flea in each lake sampled. In 2021, no mussels or spiny water flea were detected at Moose Lake.



*Eurasian watermilfoil is a non-native aquatic plant that poses a threat to aquatic habitats in Alberta because it grows in dense mats preventing light penetration through the water column, reduces oxygen levels when the dense mats decompose, and outcompetes native aquatic plants. Eurasian watermilfoil can look similar to the native Northern watermilfoil, thus genetic analysis is ideal for suspect watermilfoil species identification.*

A watermilfoil specimen was collected from Moose Lake on July 14<sup>th</sup>, and was confirmed to be the native Northern Watermilfoil.

## WATER LEVELS

*There are many factors influencing water quantity. Some of these factors include the size of the lake's drainage basin, precipitation, evaporation, water consumption, ground water influences, and the efficiency of the outlet channel structure at removing water from the lake. Requests for water quantity monitoring should go through Alberta Environment and Parks Monitoring and Science division*

Water levels at Moose Lake in 2021 were around the historical average (Figure 5). Historical data indicates that since the beginning of the record in 1950, the lake has experienced continuous dropping levels until about the early 1990s, where an appreciable increase in levels followed, and then levels decreased again in the late 1990s and early 2000s. Levels gradually increased and then jumped again in the late 2010s, and more recently are on a downward trajectory.




Figure 5. Water levels measured at Moose Lake in metres above sea level (masl) from 1950-2021. Data retrieved from Alberta Environment and Parks. Black dashed line represents historical yearly average water level.

## WEATHER & LAKE STRATIFICATION

*Air temperature will directly impact lake temperatures, and result in different temperature layers (stratification) throughout the lake, depending on its depth. Wind will also impact the degree to which a lake mixes, and how it will stratify. The amount of precipitation that falls within a lake's watershed will have important implications, depending on the context of the watershed and the amount of precipitation that has fallen. Solar radiation represents the amount of energy that reaches the earth's surface, and has implications for lake temperature & productivity.*

Moose Lake experienced a warmer, drier, windier summer with slightly more solar radiation compared to normal (Figure 6). A few warm spells prior to the July 14<sup>th</sup> sampling likely resulted in relatively high, near-surface lake temperatures. At some point during the July 14<sup>th</sup> and August 13<sup>th</sup> sampling events, the lake completely mixed, or turned over, likely during a windy spell in either late July or mid-August. This lake mixing might explain the steep increase in TP between the July and August sampling events, where anoxic bottom waters high in TP mixed with the rest of the lake water column.




Figure 6. Average air temperature (°C), accumulated precipitation (cm), and wind speed (km/h) measured from Hoselaw AGCM, as well as solar radiation (MJ/m<sup>2</sup>) measured from Dupre AGCM, with Moose Lake temperature profiles (°C) at the bottom. Black lines indicate 2021 levels, gray indicates long-term normals, and blue lines indicate sampling dates for Moose Lake over the summer. Further information about the weather data provided is available in the LakeWatch 2021 Methods report. Weather data provided by Agriculture, Forestry and Rural Economic Development, Alberta Climate Information Service (ACIS) <https://acis.alberta.ca> (retrieved April 2022).

Table 2a. Average Secchi depth and water chemistry values for Moose Lake. Historical values are given for reference.  
Number of sample trips are inconsistent between years.

Parameter	1983	1984	1985	1986	1987	1988	1990	1991	1992	1993	1994
TP (µg/L)	36	46	25	40	50	42	51	54	44	40	41
TDP (µg/L)	/	/	/	/	/	/	/	/	/	/	12
Chlorophyll- <i>a</i> (µg/L)	13.7	16.2	12.5	17.6	21.5	16	22.3	31.1	15.7	21	22.7
Secchi depth (m)	2.25	1.94	3.76	2.55	2.48	2.5	2.18	3.38	2.68	3	2.11
TKN (mg/L)	/	/	1.3	/	/	/	/	/	/	/	1.4
NO <sub>2</sub> -N and NO <sub>3</sub> -N (µg/L)	25	25	25	25	10	8	5	10	2	4	3
NH <sub>3</sub> -N (µg/L)	/	/	/	/	/	/	/	/	/	/	17
DOC (mg/L)	/	/	/	/	/	/	/	/	/	/	18
Ca (mg/L)	24	24	26	27	28	27	22	23	22	26	24
Mg (mg/L)	32	34	36	36	36	40	40	42	44	44	44
Na (mg/L)	62	64	64	66	62	74	78	74	76	82	84
K (mg/L)	12	11	12	12	12	12	12	12	13	13	14
SO <sub>4</sub> <sup>2-</sup> (mg/L)	82	84	88	92	94	102	107	106	112	117	115
Cl <sup>-</sup> (mg/L)	12	12	12	13	14	13	14	15	14	16	16
CO <sub>3</sub> (mg/L)	/	16.8	9	11.6	14.4	11.5	12.1	25.5	16	21.5	29.5
HCO <sub>3</sub> (mg/L)	/	273	289	289	283	302	294	275	300	330	330
pH	8.40	8.68	8.62	8.63	8.65	8.58	8.70	8.93	8.70	8.84	8.99
Conductivity (µS/cm)	656	641	667	678	681	715	708	706	736	780	787
Hardness (mg/L)	/	198	214	216	216	234	218	228	235	245	242
TDS (mg/L)	370	381	390	400	400	429	432	432	444	472	474
Microcystin (µg/L)	/	/	/	/	/	/	/	/	/	/	/
Total Alkalinity (mg/L CaCO <sub>3</sub> )	244	252	252	257	256	267	262	268	272	289	295

Table 2b. Average Secchi depth and water chemistry values for Moose Lake. Historical values are given for reference.  
Number of sample trips are inconsistent between years.

Parameter	1995	1996	1997	2003	2004	2005	2006	2009	2010	2011
TP (µg/L)	43	31	48	52	38	50	59	43	46	49
TDP (µg/L)	17	/	/	14	15	13	17	20	17	18
Chlorophyll- <i>a</i> (µg/L)	17.6	5.2	16.8	39.5	22.6	27.3	35.5	15.7	19	46.1
Secchi depth (m)	1.98	3.45	2.78	2.28	2.7	2.15	1.34	3.1	1.6	2.9
TKN (mg/L)	1.6	/	/	1.7	1.5	1.6	1.8	1.6	1.7	1.6
NO <sub>2</sub> -N and NO <sub>3</sub> -N (µg/L)	6	/	/	16	3	2	2	8	8	4
NH <sub>3</sub> -N (µg/L)	23	/	/	33	38	16	23	43	24	31
DOC (mg/L)	18	/	/	/	18	18	18	18	18	17
Ca (mg/L)	23	32	28	/	/	/	/	/	/	/
Mg (mg/L)	45	45	42	/	/	/	/	/	/	/
Na (mg/L)	87	84	84	111	112	114	115	117	124	114
K (mg/L)	15	14	15	12	17	20	17	20	19	20
SO <sub>4</sub> <sup>2-</sup> (mg/L)	125	124	118	149	156	151	155	165	164	156
Cl <sup>-</sup> (mg/L)	18	17	19	23	25	25	25	28	29	27
CO <sub>3</sub> (mg/L)	19	14.5	16.2	29.3	28.5	35	31.7	30.3	27.5	18
HCO <sub>3</sub> (mg/L)	321	322	314	343	350	334	346	348	358	372
pH	8.76	8.56	8.64	8.87	8.86	8.99	8.81	8.90	8.85	8.69
Conductivity (µS/cm)	793	808	776	/	934	868	947	954	964	974
Hardness (mg/L)	241	268	246	284	266	255	261	260	260	290
TDS (mg/L)	489	493	480	573	584	580	587	604	610	599
Microcystin (µg/L)	/	/	/	/	/	0.418	0.080	0.593	0.113	1.18
Total Alkalinity (mg/L CaCO <sub>3</sub> )	295	288	284	330	334	333	336	336	338	335

Table 2c. Average Secchi depth and water chemistry values for Moose Lake. Historical values are given for reference. Number of sample trips are inconsistent between years.

Parameter	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
TP (µg/L)	53	109	74	33	34	69	91	49	64	54
TDP (µg/L)	18	41	31	10	12	12	18	15	13	11
Chlorophyll- <i>a</i> (µg/L)	26.8	50	14.3	14.6	29.6	40.7	94	38	51.9	48.1
Secchi depth (m)	1.85	0.98	3.68	2.62	1.78	1.1	1.32	2.5	1.05	1.5
TKN (mg/L)	1.7	2	1.6	1.6	1.5	2.1	2.2	1.7	2	2.1
NO <sub>2</sub> -N and NO <sub>3</sub> -N (µg/L)	2	2	36	7	2	10	14	2	2	3
NH <sub>3</sub> -N (µg/L)	20	18	87	36	38	52	104	24	37	28
DOC (mg/L)	18	24	18	16	16	17	18	18	18	17
Ca (mg/L)	/	/	/	25	27	28	29	31	32	30
Mg (mg/L)	/	/	/	52	57	54	49	47	45	45
Na (mg/L)	107	116	128	110	120	110	102	99	97	94
K (mg/L)	21	24	21	18	22	21	21	20	20	20
SO <sub>4</sub> <sup>2-</sup> (mg/L)	161	151	150	168	160	148	145	138	128	140
Cl <sup>-</sup> (mg/L)	28	28	32	32	32	31	31	31	32	34
CO <sub>3</sub> (mg/L)	28.8	36.2	29.2	26.2	24.8	22.6	27	21.2	24.4	23.8
HCO <sub>3</sub> (mg/L)	358	342	353	366	368	348	338	348	308	332
pH	8.87	8.90	8.71	8.80	8.79	8.75	8.81	8.75	8.80	8.79
Conductivity (µS/cm)	993	989	996	990	994	934	918	905	875	925
Hardness (mg/L)	263	282	253	280	302	294	272	268	262	260
TDS (mg/L)	597	602	628	618	628	586	575	558	528	555
Microcystin (µg/L)	53	109	74	33	34	69	91	49	64	54
Total Alkalinity (mg/L CaCO <sub>3</sub> )	342	341	339	344	342	322	325	320	295	312



Table 3a. Concentrations of metals measured in Moose Lake. The CCME heavy metal Guidelines for the Protection of Freshwater Aquatic Life (unless otherwise indicated) are presented for reference. Note that metal sample collection method changed in 2016 from composite to single surface grab at the profile location.

Metals (Total Recoverable)	2003	2004	2005	2009	2010	2011	Guidelines
Aluminum µg/L	14.75	4.95	3.34	16.05	10.7	4.08	100 <sup>a</sup>
Antimony µg/L	0.075	0.065	0.065	0.058	0.053	0.056	/
Arsenic µg/L	1.99	2.03	2.19	2.12	2.16	2.085	5
Barium µg/L	46.1	50.2	47.8	45.4	44.9	46	/
Beryllium µg/L	0.06	0.002	0.002	0.004	0.002	0.004	100 <sup>c,d</sup>
Bismuth µg/L	0.00575	0.001	0.006	0.006	0.001	0.001	/
Boron µg/L	169.5	172	176	197	185	202	1500
Cadmium µg/L	0.030	0.007	0.005	0.005	0.005	0.004	0.37 <sup>b</sup>
Chromium µg/L	0.33	0.87	0.61	0.30	0.22	0.22	/
Cobalt µg/L	0.010	0.014	0.021	0.011	0.007	0.030	50,1000 <sup>c,d</sup>
Copper µg/L	0.56	0.75	0.61	0.49	0.26	0.50	4 <sup>b</sup>
Iron µg/L	3.25	1	37	8.05	7.65	22.8	300
Lead µg/L	0.079	0.047	0.080	0.216	0.011	0.013	7 <sup>b</sup>
Lithium µg/L	40.05	53.4	57.3	61.2	53.1	70.75	2500 <sup>d</sup>
Manganese µg/L	9.28	8.14	7.26	7.55	7.2	5.615	220 <sup>e</sup>
Molybdenum µg/L	0.590	0.846	0.705	0.598	0.556	0.628	73
Nickel µg/L	0.030	0.003	0.110	<0.005	0.003	0.163	150 <sup>b</sup>
Selenium µg/L	0.525	0.270	0.276	0.396	0.375	0.358	1
Silver µg/L	0.0025	0.003	0.001	0.002	0.002	0.008	0.25
Strontium µg/L	282.5	309	307.5	303	281	287.5	/
Thallium µg/L	0.0925	0.002	0.029	0.004	0.002	<0.002	0.8
Thorium µg/L	0.004	0.009	0.019	0.002	0.008	0.012	/
Tin µg/L	0.08	0.02	0.02	0.04	0.02	0.03	/
Titanium µg/L	0.65	0.67	0.86	1.13	0.76	0.49	/
Uranium µg/L	0.43	0.44	0.59	0.45	0.43	0.46	15
Vanadium µg/L	0.45	0.39	0.38	0.29	0.24	0.26	100 <sup>c,d</sup>
Zinc µg/L	2.98	7.9	4.335	0.722	0.498	0.68	30 <sup>f</sup>

Values represent means of total recoverable metal concentrations.

<sup>a</sup> Based on pH ≥ 6.5

<sup>b</sup> Based on 2021 avg. water hardness (as CaCO<sub>3</sub>) with CCME equation

<sup>c</sup> Based on CCME Guidelines for Agricultural use (Livestock).

<sup>d</sup> Based on CCME Guidelines for Agricultural Use (Irrigation).

<sup>e</sup> Based on CCME Manganese variable calculation ([https://ccme.ca/en/chemical/129#\\_aqf\\_fresh\\_concentration](https://ccme.ca/en/chemical/129#_aqf_fresh_concentration)), using 2021 avg. water hardness (as CaCO<sub>3</sub>) and avg. pH

<sup>f</sup> Based on 2021 avg. water hardness (as CaCO<sub>3</sub>), avg. pH, and avg. DOC with CCME equation

<sup>g</sup> Did not exceed guideline due to effect of relatively lower pH value in 2014 on Manganese variable calculation.

A forward slash (/) indicates an absence of data or guidelines

Table 3b. Concentrations of metals measured in Moose Lake. The CCME heavy metal Guidelines for the Protection of Freshwater Aquatic Life (unless otherwise indicated) are presented for reference. Note that metal sample collection method changed in 2016 from composite to single surface grab at the profile location.

Metals (Total Recoverable)	2018	2020	2021	Guidelines
Aluminum µg/L	1.3	3.7	5.9	100 <sup>a</sup>
Antimony µg/L	0.055	0.054	0.045	/
Arsenic µg/L	2.18	2.03	2	5
Barium µg/L	50.6	51.8	50.8	/
Beryllium µg/L	<0.003	<0.003	0.0015	100 <sup>c,d</sup>
Bismuth µg/L	<0.003	<0.003	0.0015	/
Boron µg/L	172	151	154	1500
Cadmium µg/L	<0.01	<0.01	0.005	0.35 <sup>b</sup>
Chromium µg/L	<0.1	<0.1	0.1	/
Cobalt µg/L	0.036	0.032	0.068	50,1000 <sup>c,d</sup>
Copper µg/L	0.3	<0.08	0.16	4 <sup>b</sup>
Iron µg/L	12.6	8.7	7.7	300
Lead µg/L	0.036	0.007	0.007	7 <sup>b</sup>
Lithium µg/L	54.1	44.6	49.7	2500 <sup>d</sup>
Manganese µg/L	11.1	21	22.6	130 <sup>e</sup>
Molybdenum µg/L	0.555	0.458	0.484	73
Nickel µg/L	0.410	0.11	0.16	150 <sup>b</sup>
Selenium µg/L	0.400	0.5	0.5	1
Silver µg/L	0.002	<0.001	0.0005	0.25
Strontium µg/L	305	293	279	/
Thallium µg/L	<0.002	<0.002	0.001	0.8
Thorium µg/L	<0.002	0.003	0.003	/
Tin µg/L	<0.06	<0.06	0.03	/
Titanium µg/L	0.69	0.57	0.5	/
Uranium µg/L	0.44	0.324	0.343	15
Vanadium µg/L	0.282	0.582	0.267	100 <sup>c,d</sup>
Zinc µg/L	5.4	0.5	0.9	30 <sup>f</sup>

Values represent means of total recoverable metal concentrations.

<sup>a</sup> Based on pH ≥ 6.5

<sup>b</sup> Based on 2021 avg. water hardness (as CaCO<sub>3</sub>) with CCME equation

<sup>c</sup> Based on CCME Guidelines for Agricultural use (Livestock).

<sup>d</sup> Based on CCME Guidelines for Agricultural Use (Irrigation).

<sup>e</sup> Based on CCME Manganese variable calculation ([https://ccme.ca/en/chemical/129#\\_aqf\\_fresh\\_concentration](https://ccme.ca/en/chemical/129#_aqf_fresh_concentration)), using 2021 avg. water hardness (as CaCO<sub>3</sub>) and avg. pH

<sup>f</sup> Based on 2021 avg. water hardness (as CaCO<sub>3</sub>), avg. pH, and avg. DOC with CCME equation

<sup>g</sup> Did not exceed guideline due to effect of relatively lower pH value in 2014 on Manganese variable calculation.

A forward slash (/) indicates an absence of data or guidelines

## LONG TERM TRENDS

Trend analysis was conducted on the parameters total phosphorus (TP), chlorophyll-*a*, total dissolved solids (TDS) and Secchi depth to look for changes over time in Moose Lake. In sum, significant increasing trends were detected for TP, chlorophyll-*a*, and TDS, and a significant decreasing trends was detected for Secchi depth. Secchi depth can be subjective and is sensitive to variation in weather - trend analysis must be interpreted with caution. Data is presented below as both line and box-and-whisker plots. Detailed methods are available in the *ALMS Guide to Trend Analysis on Alberta Lakes*.

Table 4. Summary table of trend analysis on Moose Lake data from 1983 to 2021.

Parameter	Date Range	Direction of Significant Trend
Total Phosphorus	1983-2021	Increasing
Chlorophyll- <i>a</i>	1983-2021	Increasing
Total Dissolved Solids	1983-2021	Increasing
Secchi Depth	1983-2021	Decreasing

## Definitions:


**Median:** the value in a range of ordered numbers that falls in the middle.

**Trend:** a general direction in which something is changing.

**Monotonic trend:** a gradual change in a single direction.

**Statistically significant:** The likelihood that a relationship between variables is caused by something other than random chance. This is indicated by a *p*-value of  $<0.05$ . **Variability:** the extent by which data is inconsistent or scattered.

**Box and Whisker Plot:** a box-and-whisker plot, or boxplot, is a way of displaying all of our annual data. The median splits the data in half. The 75<sup>th</sup> percentile is the upper quartile of the data, and the 25<sup>th</sup> percentile is the lower quartile of the data. The top and bottom points are the largest and smallest observations.



## Total Phosphorus (TP)

Trend analysis of TP over time showed that it is significantly increasing in Moose Lake since 1983 (Tau = 0.19,  $p = 0.0048$ ; Table 5).




Figure 6. Monthly total phosphorus (TP) concentrations measured between June and September on sampling dates between 1983 and 2021 ( $n = 114$ ). The value closest to the 15<sup>th</sup> day of the month was chosen to represent the monthly value in cases with multiple monthly samples.

## Chlorophyll-a

Chlorophyll-a has been significantly increasing in Moose Lake since 1983 (Tau = 0.21,  $p = 0.0014$ ; Table 5). TP and chlorophyll-a were significantly correlated ( $r = 0.53$ ,  $p = 2.02 \times 10^{-9}$ ).




Figure 7. Monthly chlorophyll-a concentrations measured between June and September on sampling dates between 1983 and 2021 ( $n = 114$ ). The value closest to the 15<sup>th</sup> day of the month was chosen to represent the monthly value in cases with multiple monthly samples. Line graph is overlain by TP concentrations.



## Total Dissolved Solids (TDS)

Trend analysis showed a significant increasing trend in TDS between 1983 and 2021 ( $\text{Tau} = 0.48$ ,  $p = <0.001$ ) in Moose Lake. Despite the overall trend, TDS began to decrease in 2016 (Figure 8), which could be related to changes in water levels at this time (Figure 5).




Figure 8. Monthly TDS values measured between June and September on sampling dates between 1983 and 2021 ( $n = 81$ ). The value closest to the 15<sup>th</sup> day of the month was chosen to represent the monthly value in cases with multiple monthly samples.

Due to the significant increasing trend of TDS in Moose Lake, exploring the specific major ions which may be driving this trend is important to determine. Trend analysis of major ions at Moose Lake indicates that sulphate, alkalinity (bicarbonate, carbonate), and sodium are likely the key parameters that are driving the increase in TDS (Figure 9). These three parameters also follow the TDS trajectory the most closely over time, but significant increases in potassium and chloride are likely also contributing to increases in TDS over time. Interestingly, neither potassium nor chloride display decreases in recent years unlike TDS, sulphate, alkalinity and sodium, but rather display slight stabilization and continued increases, respectively.




Figure 9. Concentrations of TDS (top left, blue panel), major ions (sodium = Na<sup>+</sup>, magnesium = Mg<sup>2+</sup>, potassium = K<sup>+</sup>, calcium = Ca<sup>2+</sup>, chloride = Cl<sup>-</sup>, sulphate = SO<sub>4</sub><sup>2-</sup>), and total alkalinity (Alk., as mg/L CaCO<sub>3</sub>) measured monthly between June and September on sampling dates between 1983 and 2021. Also represented is the monotonic trend results for each parameter; test used (MK = Mann Kendall, SK = Seasonal Kendall), significance of test ( $p$ ; assessed as significance when  $p < 0.05$ , marked with “\*” if significant), and the slope of the trend. Test selection follows method outline in the *ALMS Guide to Trend Analysis on Alberta Lakes*. Note that some ions had insufficient data (\*I.D.\*) therefore trends were not calculated. The value closest to the 15<sup>th</sup> day of the month was chosen to represent the monthly value in cases with multiple monthly samples.

## Secchi Depth

Trend analysis of Secchi depth over time indicates that it is significantly decreasing in Moose Lake since 1983 (Tau = -0.20,  $p = 0.0028$ ; Table 5).




Figure 9. Monthly Secchi depth values measured between June and September on sampling dates between 1983 and 2021 ( $n = 116$ ). The value closest to the 15<sup>th</sup> day of the month was chosen to represent the monthly value in cases with multiple monthly samples.

Table 5. Results of trend tests using total phosphorus (TP), chlorophyll-*a*, total dissolved solids (TDS) and Secchi depth data from June to September, for sampled years from 1983-2021 on Moose Lake data.

Definition	Unit	Total Phosphorus (TP)	Chlorophyll-a	Total Dissolved Solids (TDS)	Secchi Depth
Statistical Method	-	Seasonal Kendall	Seasonal Kendall	Seasonal Kendall	Seasonal Kendall
The strength and direction (+ or -) of the trend between -1 and 1	Tau	0.19	0.21	0.48	-0.20
The extent of the trend	Slope (units per year)	0.32	0.29	5.15	-0.02
The statistic used to find significance of the trend	Z	2.82	3.19	5.96	-2.98
Number of samples included	n	114	114	81	116
The significance of the trend	<i>p</i>	0.0048*	0.0014*	$2.52 \times 10^{-9}$ *	0.0028*

\**p* < 0.05 is significant within 95%