



*The Alberta Lake Management Society  
Volunteer Lake monitoring report*

# Laurier Lake



## 2007 Report

*Completed with support from:*



**Alberta Lake Management Society**

**Address:** PO Box 4283, Edmonton,  
Alberta, T6E 4T3

**Phone:** 780-702-ALMS

**E-mail:** info@alms.ca

*Water is integral to supporting and maintaining life on this planet as it moderates the climate, creates growth and shapes the living substance of all of Earth's creatures. It is the tide of life itself, the sacred source.* David Suzuki (1997). The Sacred Balance.

## Alberta Lake Management Society's Lakewatch Program

Lakewatch has several important objectives, one of which is to collect and interpret water quality on Alberta Lakes. Equally important is educating lake users about their aquatic environment, encouraging public involvement in lake management, and facilitating cooperation and partnerships between government, industry, the scientific community and lake users. Lakewatch Reports are designed to summarize basic lake data in understandable terms for a lay audience and are not meant to be a complete synopsis of information about specific lakes. Additional information is available for many lakes that have been included in Lakewatch and readers requiring more information are encouraged to seek these sources.

ALMS would like to thank all who express interest in Alberta's aquatic environments and particularly those who have participated in the Lakewatch program. These people prove that ecological apathy can be overcome and give us hope that our water resources will not be the limiting factor in the health of our environment.

### Acknowledgements

The Lakewatch program is made possible through the dedication of volunteers and the Lakewatch Chairs, Théo Charette and Ron Zurawell. We thank Bev Smith for her time and effort in the collection of field data and the use of her watercraft during 2007. Numerous Alberta Environment staff also contributed to successful completion of the 2007 program. We would like to thank Jill Anderson and Wendy Markowski who were summer interns with ALMS in 2007. Project Technical Coordinator, Megan McLean was instrumental in planning and organizing the field program. Technologists, Mike Bilyk, Brian Jackson and John Willis were involved in the logistics planning and training aspects of the program. Doreen LeClair was responsible for data management. Theo Charette (ALMS Director) was responsible for program administration and planning. Théo Charette, Ron Zurawell (Limnologist, AENV), and Lori Nuefeld prepared the original report, which was updated by Heather Powell for the 2007 report. Alberta Environment, Lakeland Industry and Community Association (LICA) financially supported the Lakewatch program.


# Laurier Lake

Laurier Lake is one of four beautiful lakes (**Figure 1**) that were left behind 10 000 years ago when glaciers carved a setting of hummocky terrain of kettles, eskers and lake basins. Archeological evidence indicates the area was inhabited at least 7 000 years ago. The first Europeans came through the area in 1754 by way of the nearby North Saskatchewan River.



**Figure 1.** Laurier Lake, Alberta. Photo: L. Kowalchuk, ALMS

The Whitney Lakes Provincial Park adjacent to Laurier Lake (**Figure 2**) was established in 1982. It boasts a diverse setting of jack pine (*Pinus banksiana*) meadows, aspen (*Populus spp.*) groves, willow (*Salix spp.*) thickets, marshes, fens, and mixed wood forests. As many as 148 bird species have been observed in the park with an excellent viewing point on the west side of Laurier Lake (SRD 2002). The land surrounding Laurier Lake includes a mixture of recreational cottage development, cleared agricultural land and natural deciduous forest. Protected Crown land makes up the north shore of the lake the remainder of the shoreline is privately owned (Mills 1988). The lake is enjoyed for recreational activities such as hiking, wildlife viewing and water-based recreation.



**Figure 2.** Bathymetry of Laurier Lake, Alberta. Contours are at 5 ft. intervals


Popular activities include: wind surfing, water-skiing, sailing, swimming and fishing. Yellow perch (*Perca flavescens*), walleye (*Stixostedion vitreum*) and northern pike (*Esox lucius*) are the sport fish of Laurier Lake. Fish stocking occurred in 1953. Sport and forage fish were transferred from Moose Lake to Laurier Lake. The lake has not been managed for commercial or domestic fishing.

Laurier Lake has a surface area of 6.42 km<sup>2</sup> with a maximum 2004 depth of 6.6 m (**Figure 2**). The lake has been both mesotrophic and eutrophic. Its location and surrounding topography make Laurier open to prevailing winds. These winds mix the water column and Laurier usually does not thermally stratify throughout most of the

summer. Mixing also allows nutrients and organic material to remain suspended in the water column making the lake naturally fertile. Algal blooms are known to occur during summer months due to the lakes natural fertility. Detailed studies on phytoplankton have not been completed for the lake. Common emergent plants that fringe the lake are bulrushes (*Scirpus* spp.), cattails (*Typha latifolia*) and sedges (*Carex* spp.).

### *Water Levels*

Laurier Lake shares a 92-km<sup>2</sup> drainage area with Ross, Borden and Whitney Lakes. One intermittent and three permanent streams feed the lake. The outflow, on the northwest end, drains into Borden Lake and subsequently to the North Saskatchewan River (Mills 1988).



**Figure 3.** Historical water level elevation (meters above sea level (asl)) in Laurier Lake, Alberta, 1968-2007.

Water levels in Laurier Lake have been monitored since 1968. From the historical records, the water level was at a maximum in 1974 but dropped almost 3 m to a minimum recorded level in 2004 (**Figure 3**). Water levels have been slowly dropping for the last 2 decades, with a slight increase from 2005-2007. The average elevation is 565.7 m above sea level. Alberta experienced a relatively wet year in 1997 that restored water levels in many lakes. In Laurier Lake, the wet year of 1997 temporarily halted water level declines. The reprieve was short-lived and water levels declined again after 1997.

## Results


### *Water Temperature and Dissolved Oxygen*

*Water temperature and dissolved oxygen profiles in the water column can provide information on water quality and fish habitat. Please refer to the end of this report for descriptions of technical terms.*

Laurier Lake is a polymictic lake, which mixes many times throughout the summer. Because of frequent mixing the water temperature and dissolved oxygen concentration is relatively the same throughout the lake (**Figure 4**). Water temperature peaked in mid

July and a slight thermal stratification was noted at 2m. Laurier Lake mixed prior to the next sample date (8 August) and thermal stratification was no longer evident.

Dissolved oxygen (DO) concentrations were the same at all depths in June 2007 (**Figure 4**). In mid-July, DO increased slightly in surface water and at 3m depth. The peak DO concentration at 3m indicates a phytoplankton layer that is concentrated at the thermocline. DO concentrations declined at 4 m in July, at 5 m in August and 6 m in September. DO approached zero (e.g. anoxic) near the lake bed in September. The pattern of DO concentrations in July-September reflects microbial decomposition, which consumes oxygen. The oxygen levels in surface layers of Laurier Lake were within the acceptable range for surface water quality, according to Alberta Environment guidelines ( $DO \geq 5.0$  mg/L).



**Figure 4.** Water temperature (°C) and dissolved oxygen (mg/L) profiles for Laurier Lake during the summer of 2007.

#### *Water clarity and Secchi Depth*


*Suspended material, both living and dead, as well as some coloured dissolved compounds in the water column influence water clarity. The most widely used measure of lake water clarity is the Secchi depth. After ice and snowmelt, a lake can have low clarity due to spring runoff and suspended sediments in the lake. Lake water usually clears in the spring but then becomes less clear as algae grow through the summer.*

Laurier Lake is a shallow, polymictic lake and as such is relatively clear, with average Secchi disk depth = 2.4 m. During the summer 2007, light penetrated to an average 37% of the total lake depth, thus algae could grow in the upper 1/3 of the lake (**Table 1**). Water clarity peaked on 11 June (Secchi disk depth = 3.5 m) early in the growing season

and before the lake mixed. Water clarity declined in September (Secchi disk depth = 1.5 m) after Laurier Lake mixed (**Figure 4**). The decrease in water clarity with over-turn is the result of particle re-suspension and is a common characteristic of polymictic lakes. As particles settle after mixing, water clarity increases.

### Water chemistry

Based on lake water characteristics, Laurier Lake is mesotrophic to slightly eutrophic (see *A Brief Introductory to Limnology* at the end of this report). Total phosphorous concentration peaked in June (TP = 51 µg/L) and was lowest in mid-July (**Figure 5**). Phosphorous was within the mesotrophic range. Nitrogen concentration was relatively stable throughout the summer (mean TN = 2.2 mg/L, **Figure 5**) and was within the



**Figure 5.** Total phosphorus, total nitrogen, and chlorophyll *a* (a measure of algae biomass) concentrations for Laurier Lake during the summer of 2007.

hypereutrophic range. Algae biomass (chlorophyll *a*) in Laurier Lake peaked in early August (maximum chl *a* = 6.8; mean chl *a* = 4.3, **Figure 5**) and was within the mesotrophic range. Algal growth was phosphorous limited in Laurier Lake in 2007.

Nutrient concentrations remained relatively stable on an annual mean basis. There is no evidence to suggest increased nutrient loading from cottages or other land use activities around Laurier Lake.

Compared to other Alberta Lakes, algal biomass in Laurier Lake was moderate and relatively stable over the course of the summer. Algal blooms were not observed in 2007 or the last reporting date 2004.

Laurier Lake is well buffered. In 2007, lake pH = 9.1, which is well above that of pure water (i.e., pH 7). Laurier Lake is a hardwater, slightly saline lake. Ion concentrations were relatively high 2007 (**Table 1**). Dominate ions were bicarbonate, sodium, and sulfate. Magnesium was not measured in 2007. Mineral ions such as sulfate are supplied by weathering in the watershed and from groundwater inflows. The change in ion concentration of Laurier Lake suggests a fundamental change in its hydrology. Low rainfall has likely reduced the contribution of runoff to Laurier Lake. With a reduction in runoff and particularly through flow (lateral flow through surface soil), groundwater has likely become a more important source of water to Laurier Lake.

The average concentrations of various heavy metals (as total recoverable concentrations) in Laurier Lake did not surpass provincial and federal Water Quality Guidelines for the

Parameter	1978 JUNE	1980 AUG	1987 AUG	1997	1998	2000	1999 FEB	2001 SEP	2002	2003	2004	2007
TP (µg/L)	-	-	-	32	48	37	16	-	36	27	40	41.6
TDP (µg/L)	-	-	-	-	-	-	22	-	15	15	18	22
Chla (µg/L)	-	-	-	5.3	8.9	5.5	1.1	-	5.8	2.6	5.0	4.3
Secchi (m)	-	1.3	1.2	4.6	1.3	1.8	1.7	2.3	2.5	4.4	3.2	2.4
TKN (mg/L)	-	-	-	-	-	-	-	-	2.5	2.6	2.7	2.2
NO <sub>2+3</sub> (µg/L)	<50	50	<1	-	-	-	6	-	3.8	2.11	7.7	<7
NH <sub>4</sub> (µg/L)	-	-	-	-	-	-	25	-	23	41	76	46.2
Dissolved organic C (mg/L)	-	-	-	-	-	-	-	-	-	-	-	37.9
Ca (mg/L)	23	27	19	20	21	13	18	-	12	10	10.5	14.6
Mg (mg/L)	48	54	52	73	81	83	86	-	99	106	107	-
Na (mg/L)	49	45	59	86	92	98	103	-	77	128	129	122.7
K (mg/L)	14	14	17	24	25	25	27	-	26	31	34	32.8
SO <sub>4</sub> (mg/L)	36	40	41	62	66	73	74	-	94	99	105	111.7
Cl (mg/L)	5	6	9	12	13	15	17	-	12	18	20	19.5
CO <sub>3</sub> (mg/L)	-	-	-	39	62	66	-	-	102	112	84	86
HCO <sub>3</sub> (mg/L)	-	-	-	493	468	469	-	-	515	522	603	535.7
pH	-	-	-	8.8	8.9	8.0	-	-	9.2	9.2	9.1	9.1
Conductivity (µS/cm)	-	-	-	-	-	-	-	-	-	-	-	1163
TDS (mg/L)	-	-	-	562	598	602	-	-	-	764	-	-
Total Alkalinity (mg/L CaCO <sub>3</sub> )	310	329	360	470	488	493	562	-	592	615	634	583

Protection of Aquatic Life (**Appendix 1**).

**Table 1.** Historical mean water chemistry values for Laurier Lake, Alberta. Average values for summer 2007 and previous years. Values are summer averages, unless noted.

**Note.** TP = total phosphorous, TDP = total dissolved phosphorus, Chla = chlorophyll *a*, TKN = total Kjeldahl nitrogen, NO<sub>2+3</sub> = nitrate+nitrite, NH<sub>4</sub> = ammonium, DOC = dissolved organic carbon, Ca = calcium, Mg = magnesium, Na = sodium, K = potassium, SO<sub>4</sub> = sulphate, Cl = chloride, CO<sub>3</sub> = carbonate, HCO<sub>3</sub> = bicarbonate, TDS = total dissolved solids.

## References

Mills 1988

Mitchell, P. and E. Prepas, eds. 1990. *Atlas of Alberta Lakes*. University of Alberta Press.



SRD 2002

## Appendix 1

Mean concentrations of metals in Laurier Lake for summer 2007. The CCME heavy metal Guidelines for the Protection of Freshwater Aquatic Life (unless otherwise indicated) are presented for reference.

<b>Metals</b>	<b>2007</b>	<b>Guidelines</b>
ALUMINUM µg/L	29.4	100 <sup>a</sup>
ANTIMONY µg/L	0.137	6 <sup>e</sup>
ARSENIC µg/L	2.6	5
BARIUM µg/L	20.2	1000 <sup>e</sup>
BERYLLIUM µg/L	<0.003	100 <sup>d,f</sup>
BISMUTH µg/L	<0.005	-
BORON µg/L	175.5	5000 <sup>e,f</sup>
CADMIUM µg/L	0.008	0.085 <sup>b</sup>
CHROMIUM µg/L	0.611	-
COBALT µg/L	0.099	1000 <sup>f</sup>
COPPER µg/L	0.613	4 <sup>c</sup>
IRON µg/L	37.1	300
LEAD µg/L	0.057	7 <sup>c</sup>
LITHIUM µg/L	102.9	2500 <sup>g</sup>
MANGANESE µg/L	5.15	200 <sup>g</sup>
MOLYBDENUM µg/L	0.661	73 <sup>d</sup>
NICKEL µg/L	0.275	150 <sup>c</sup>
SELENIUM µg/L	0.547	1
SILVER µg/L	<0.003	0.1
STRONTIUM µg/L	84.5	-
THALLIUM µg/L	<0.001	0.8
THORIUM µg/L	<0.01	-
TIN µg/L	<0.06	-
TITANIUM µg/L	1.24	-
URANIUM µg/L	0.811	100 <sup>e</sup>
VANADIUM µg/L	0.742	100 <sup>f,g</sup>
ZINC µg/L	1.53	30
FLUORIDE mg/L	-	1.5

With the exception of fluoride (which reflects the mean concentration of dissolved fluoride only), values represent means of total recoverable metal concentrations.

<sup>a</sup> Based on pH ≥ 6.5; calcium ion concentration [Ca<sup>+2</sup>] ≥ 4 mg/L; and dissolved organic carbon concentration [DOC] ≥ 2 mg/L.

<sup>b</sup> Based on water Hardness of 300 mg/L (as CaCO<sub>3</sub>).

<sup>c</sup> Based on water Hardness > 180 mg/L (as CaCO<sub>3</sub>).

<sup>d</sup> CCME interim value.

<sup>e</sup> Based of Canadian Drinking Water Quality guideline values.

<sup>f</sup> Based of CCME Guidelines for Agricultural Use (Livestock Watering).

<sup>g</sup> Based of CCME Guidelines for Agricultural Use (Irrigation).


# A brief introduction to Limnology

## *Indicators of water quality*

The goal of **Lakewatch** is to collect water samples necessary to determine the water quality of lakes. Though not all encompassing, the variables measured in **Lakewatch** are sensitive to human activities in watersheds that may cause impacts to water quality. For example, nutrients such as phosphorus and nitrogen are important determinants of lake productivity. The concentrations of these nutrients in a lake are affected (typically elevated) by land use changes such as increased crop production or livestock grazing. Elevated nutrient concentrations can cause increases in undesirable algae blooms resulting in low dissolved oxygen concentrations, degraded fish habitat and production of noxious odors. Large increases in nutrients over time may also indicate sewage inputs, which in turn, may result in other human health concerns such as harmful bacteria or protozoans (e.g. *Cryptosporidium*).

## *Temperature and mixing*

Water temperature in a lake dictates the behavior of many chemical parameters responsible for water quality (Figure 6). Heat is transferred to a lake at its surface and slowly moves downward depending on water circulation in the lake. Lakes with a large surface area or a small volume tend to have greater mixing due to wind. In deeper lakes, circulation is not strong enough to move warm water to depths typically greater than 4 or 5 m and as a result cooler denser water remains at the bottom of the lake. As the difference in temperature between warm surface and cold deeper water increases, two distinct layers are formed. Limnologists call these layers of water the **epilimnion** at the surface and the **hypolimnion** at the bottom. A transition layer known as the metalimnion, which contains the effective wall separating top and bottom waters called a thermocline, separates the layers. A thermocline typically occurs when water temperature changes by more than one degree within one-meter depth. The hypolimnion and epilimnion do not mix, nor do elements such as oxygen supplied at the surface move downward into the hypolimnion. In the fall, surface waters begin to cool and eventually reach the same temperature as hypolimnetic water. At this point, the water mixes from top to bottom in what is called a **turnover** event. Surface water cools further as ice forms and again a thermocline develops this time with 4° C water at the bottom and 0° C water on the top.



**Figure 6: Difference in the circulation of the water column depending on thermal stratification.**

In spring, another turnover event occurs when surface waters warm to 4° C. Lakes with this mixing pattern of two stratification periods and two turnover events are called **dimictic** lakes. In shallower lakes, the water column may mix from top to bottom most of the ice-free season with occasional stratification during periods of calm warm conditions. Lakes that mix frequently are termed **polymictic** lakes. In our cold climate, many shallow lakes are **cold monomictic** meaning a thermocline develops every winter, there is one turnover event in spring but the remainder of the ice-free season the lake is polymictic.

## *Dissolved Oxygen*

Oxygen enters a lake at the lake surface and throughout the water column when produced by photosynthesizing plants, including algae, in the lake. Oxygen is consumed within the lake by respiration of living organisms and decomposition of organic material in the lake sediments. In lakes that stratify (see temperature above), oxygen that dissolves into the lake at the surface cannot mix downward into the hypolimnion. At the same time, oxygen is depleted in the hypolimnion by decomposition. The result is that

the hypolimnion of a lake can become **anoxic**, meaning it contains little or no dissolved oxygen. When a lake is frozen, the entire water column can become anoxic because the surface is sealed off from the atmosphere. Winter anoxic conditions can result in a fish-kill, which is particularly common during harsh winters with extended ice-cover. Alberta Surface Water Quality Guidelines suggest dissolved oxygen concentrations (in the epilimnion) must not decline below 5 mg/L and should not average less than 6.5 mg/L over a seven-day period. However, the guidelines also require that dissolved oxygen concentrations remain above 9.5 mg/L in areas where early life stages of aquatic biota, particularly fish, are present.

### *General Water Chemistry*

Water in lakes always contains substances that have been transported by rain and snow or have entered the lake in groundwater and inflow streams. These substances may be dissolved in the water or suspended as particles. Some of these substances are familiar minerals, such as sodium and chloride, which when combined form table salt, but when dissolved in water separate into the two electrically charged components called **ions**. Most dissolved substances in water are in ionic forms and are held in solution due to the polar nature of the water molecule. **Hydrophobic** (water-fearing) compounds such as oils contain little or no ionic character, are non-polar and for this reason do not readily dissolve in water. Although hydrophobic compounds do not readily dissolve, they can still be transported to lakes by flowing water. Within individual lakes, ion concentrations vary from year to year depending on the amount and mineral content of the water entering the lake. This mineral content can be influenced by the amount of precipitation and other climate variables as well as human activities such as fertilizer and road salt application.

### *Phosphorus and Nitrogen*

Phosphorus and nitrogen are important nutrients limiting the growth of algae in Alberta lakes. While nitrogen usually limits agricultural plants, phosphorus is usually in shortest supply in lakes. Even a slight increase of phosphorus in a lake can, given the right conditions, promote algal blooms causing the water to turn green in the summer and impair recreational uses. When pollution originating from livestock manure and human sewage enters lakes, not only are the concentrations of phosphorus and nitrogen increased but nitrogen can become a limiting nutrient, which is thought to cause blooms of toxic algae belonging to the cyanobacteria. Not all cyanobacteria are toxic, however, the blooms can form decomposing mats that smell and impair dissolved oxygen concentrations in the lake.

### *Chlorophyll-a*

Chlorophyll-*a* is a photosynthetic pigment that green plants, including algae, possess enabling them to convert the sun's energy to living material. Chlorophyll-*a* can be easily extracted from algae in the laboratory. Consequently, chlorophyll-*a* is a good estimate of the amount of algae in the water. Larger aquatic plants, known as macrophytes, rather than algae, dominate some highly productive lakes. In these lakes, chlorophyll-*a* and nutrient values taken from water samples do not include productivity from large aquatic plants. As a result, lakes like Chestermere, which are dominated by macrophytes, can exist at a lower trophic state than if macrophyte biomass was included. Unfortunately, the productivity and nutrient cycling contributions of macrophytes are difficult to sample accurately and are therefore not typically included in trophic state indices.

### *Secchi Disk Depth*


Lakes that are clear are more attractive for recreation, whereas those that are turbid or murky are considered by lake users to have poor water quality. Secchi disk depth is the oldest, simplest, and quickest quantitative measure of water clarity. A Secchi disk is a black and white disk that is lowered down through the water column until it can no longer be seen. Secchi disk depth is the midpoint between the depth at which it disappears when lowered and reappears when it is pulled up again. The Secchi disk depth in lakes with high algal biomass will generally be shallow. However, algae do not only affect Secchi disk depth.

High concentrations of suspended sediments, particularly fine clays or glacial till, are common in plains or mountain reservoirs of Alberta. Mountain reservoirs may have exceedingly shallow Secchi disk depths despite low algal growth and nutrient concentrations.

The euphotic zone, calculated as twice the Secchi disk depth, is the portion of the water column that has sufficient light for aquatic plants to grow. Murky waters, with shallow Secchi depths, can prevent aquatic plants from growing on the lake bottom. Aquatic plants are important because they ensure clear lake water by reducing shoreline erosion and stabilizing lake bottom sediments. Many lakes in Alberta are shallow and have bottom sediments with high concentrations of nutrients. Without aquatic plants, water quality may decline in these lakes due to murky, sediment-laden water and excessive algal blooms. Maintaining aquatic plants in certain areas of a lake is often essential for ensuring good water clarity and a healthy lake as many organisms, like aquatic invertebrates and fish, depend on aquatic plants for food and shelter.

### Trophic state

Trophic state is a classification system for lakes that depends on fertility and is a useful index for rating and comparing lakes. From low to high nutrient and algal biomass (as chlorophyll-*a*) concentrations, the trophic states are: **oligotrophic**, **mesotrophic**, **eutrophic** and **hypereutrophic**. The nutrient and algal biomass concentrations that define these categories are shown in table 2 and a graph of Alberta lakes compared by trophic state can be found on the ALMS website. A majority of lakes in Alberta are meso- to eutrophic because they naturally contain high nutrient concentrations due to our deep fertile soils. Thus, lakes in Alberta are susceptible to human impacts because they are already nutrient-rich; any further nutrient increases can bring about undesirable conditions illustrated in Figure. 7.



**Figure 7: Suggested changes in various lake characteristics with eutrophication. From “Ecological Effects of Wastewater”, 1980.**

**Table 2: Trophic status based on lake water characteristics**

Trophic state	Total Phosphorus (µg/L)	Total Nitrogen (µg/L)	Chlorophyll a (µg/L)	Secchi Depth (m)
Oligotrophic	< 10	< 350	< 3.5	> 4
Mesotrophic	10 - 30	350 - 650	3.5 - 9	4 - 2
Eutrophic	30 - 100	650 - 1200	9 - 25	2 - 1
Hypereutrophic	> 100	> 1200	> 25	< 1

Note: These values are from a detailed study of global lakes reported in Nurnberg 1996. Alberta Environment uses slightly different values for TP and CHL based on those of the OECD reported by Vollenweider (1982). The AENV and OECD cutoffs for TP are 10, 35 and 100; for CHL are 3, 8 and 25. AENV does not have TN or Secchi depth criteria. The corresponding OECD exists for Secchi depth and the cutoffs are 6, 3 and 1.5 m.