



*The Alberta Lake Management Society
Volunteer Lake Monitoring Program*

Marie Lake

2009 Report

Completed with support from:



Alberta Lake Management Society

Address: P.O. Box 4283
Edmonton, AB T6E4T3
Phone: 780-702-ALMS
E-mail: info@alms.ca

Water is integral to supporting and maintaining life on this planet as it moderates the climate, creates growth and shapes the living substance of all of Earth's creatures. It is the tide of life itself, the sacred source.

David Suzuki (1997) The Sacred Balance

Alberta Lake Management Society's Lakewatch Program

Lakewatch has several important objectives, one of which is to collect and interpret water quality data on Alberta Lakes. Equally important is educating lake users about their aquatic environment, encouraging public involvement in lake management, and facilitating cooperation and partnerships between government, industry, the scientific community and lake users. Lakewatch Reports are designed to summarize basic lake data in understandable terms for a lay audience and are not meant to be a complete synopsis of information about specific lakes. Additional information is available for many lakes that have been included in Lakewatch and readers requiring more information are encouraged to seek these sources.

ALMS would like to thank all who express interest in Alberta's aquatic environments and particularly those who have participated in the Lakewatch program. These people prove that ecological apathy can be overcome and give us hope that our water resources will not be the limiting factor in the health of our environment.

Acknowledgements

The Lakewatch program is made possible through the dedication of its volunteers and Lakewatch Chairs, Al Sosiak and Ron Zurawell. We would like to thank Don Savard, Steve Vinek, Julian & Harriett Nowicki, Robina Sobey, and Roy Bibeau for their efforts in collecting data in 2009. We would also like to thank Noemie Jenni and Cristen Symes who were summer interns with ALMS in 2009. Project Technical Coordinator, Jill Anderson was instrumental in planning and organizing the field program. Technologists, Shelley Manchur, Mike Bilyk, Brian Jackson and John Willis were involved in the logistics planning and training aspects of the program. Doreen LeClair and Chris Rickard were responsible for data management. Théo Charette (ALMS President) and Jill Anderson (Program Manager) were responsible for program administration and planning. Théo Charette, Ron Zurawell (Limnologist, AENV), and Lori Neufeld prepared the original report, which was updated by Sarah Lord for 2009. Alberta Environment and the Beaver River Watershed Alliance (BRWA) were major sponsors of the Lakewatch program.

Marie Lake

Marie Lake (**Figure 1**) is located in the Beaver River Drainage Basin. It lies about 26 km north of the Town of Cold Lake. Marie Lake is named after the Cree word *Methae* or *Merai*, meaning 'a fish', and may refer specifically to the burbot (*Lota lota*) prevalent throughout most of Alberta.




Figure 1. Marie Lake, Alberta, July 2004. Theo Charette, Alberta Environment.

The Cree arrived in the late eighteenth century, displacing Beaver, Blackfoot, and Slavey tribes that were common in the area. The Cree arrived during the growth of the fur trade along a popular route from Waterhen, Saskatchewan by way of the Beaver River (Mitchell and Prepas 1990).

Marie Lake is over 26 m deep (**Figure 2**) with a slow flushing rate (a residence time of 14.5 years). It is mesotrophic and has a small littoral zone for its surface area of 36 km². The shoreline is primarily sandy with macrophytes limited to a couple areas. A large macrophyte bed is located along the west shore stretching toward the north, and another lies on the western edge of the south bay. Macrophyte beds are dominated by bulrush (*Scirpus* spp.),

pondweed (*Potamogeton* spp.), and northern watermilfoil (*Myriophyllum exalbescens*)




Figure 2. Bathymetry of Marie Lake, Alberta. Contour intervals represent 3 m depth intervals.

(Mitchell and Prepas 1990). The low productivity of the shoreline does not provide suitable habitat for semi-aquatic wildlife; however, the macrophyte beds present are very important for maintaining a productive fishery.

Oil and gas extraction, and agriculture are the primary industries near Cold Lake. Orthic grey luvisols are found in the upper- to mid-slopes of the Marie Lake watershed, while poorly drained organic soils associated with moist areas (known as fibrisols and meisols) are found in depressions and flat areas. Peatlands, bogs, fens, and marshes make up an important component of the Marie Lake watershed (Mitchell and Prepas 1990). The Marie Lake Air & Watershed Society is spearheading the protection and enhancement of the Marie Lake watershed.

Marie Lake remains largely undeveloped compared with other Alberta lakes of similar size, with individual private cottages along the southern shore and a subdivision on the eastern shore on private land. Additional private property and a campground are located at Shelter Bay on the north shore. The most popular recreational activities include swimming, fishing, camping, sightseeing, and relaxing. Also popular are power boating, waterskiing, canoeing, hiking, and photography. Marie Lake currently supports a domestic fishery; commercial licences have not been issued in the last two decades. Sport fish include lake whitefish, tullibee, walleye, ling, suckers, northern pike, and yellow perch (Mitchell and Prepas 1990).

The water in Marie Lake is usually clear with little algal growth. Phytoplankton (algae) have been identified in previous studies: six species of *Cyanophyta* (blue-green algae) were present but comprised a low proportion of total algal biomass throughout the open water season. *Bacillariophyta* (diatoms), *Chrysophyta* (golden-brown algae), and *Pyrrhophyta* (Dinoflagellates) were abundant in late summer through fall.

Results

Water Levels

Water levels in Marie Lake have been monitored since 1980 (**Figure 3**). Water levels declined to a minimum 573.1 m in 1992; and subsequently increased to a maximum 574.4 m in 1997, a difference of 1.35 m. Water levels in Marie Lake fluctuate in response to changes in weather and runoff, but no significant long-term changes in water level are apparent.

The water management plan implemented in 1985 indicates




Figure 3. Historical water levels (m asl) in Marie Lake, Alberta 1980 – 2009.

water allocation from Marie Lake may reach $0.425 \times 10^6 \text{ m}^3/\text{year}$ (Mitchell and Prepas 1990). However, there are currently no industrial water withdrawals taken from Marie Lake directly. Local residents are concerned that sustained industrial withdrawals from groundwater and the potential for reduced surface water levels in the region.

Water Temperature and Dissolved Oxygen

Water temperature and dissolved oxygen profiles in the water column can provide information on water quality and fish habitat. Please refer to the end of this report for descriptions of technical terms.

Marie Lake was thermally stratified in the summer of 2009 (**Figure 4**). Surface water temperature was 9.1°C on 7 June and increased to a seasonal maximum of 20.3°C by 27 July. Surface water temperature cooled to 18.3°C on 17 August and 17.9°C on 30 August. Field sampling of the water column in June only went to 11 m, so the full depth profile of the lake was not obtained. A strong thermocline was observed on subsequent sampling dates in July and August, with the upper layer of warm water (e.g. epilimnion) extending to 7 m in July and 9 m in August. The strong thermocline present on these sampling dates indicates that waters in Marie Lake did not turnover during the summer sampling period.




Figure 4. Water temperature ($^\circ\text{C}$) and dissolved oxygen (mg/L) profiles for Marie Lake during the summer of 2009.

On all sampling dates, the oxygen levels in the surface layer of Marie Lake were within the acceptable range for surface water quality, according to Alberta Environment guidelines ($\text{DO} \geq 5.0 \text{ mg/L}$) (**Figure 4**). In June, dissolved oxygen (DO) concentrations were $>10 \text{ mg/L}$ at the surface but a complete depth profile was not available. By 27 July, DO concentrations in the bottom layers of water (e.g. hypolimnion) had begun to decrease below 7 m depth. On 17 August and 30 of August a rapid decline in DO concentration was observed at 9 m depth, as DO began to decrease and neared zero (e.g. anoxic) at the lake

bed. This DO pattern is typical for a thermally stratified lake. Deep-water anoxia is common, as bacterial decomposition of organic matter in lake sediments consumes oxygen at the lakebed.

Water Clarity and Secchi Depth

Water clarity is influenced by suspended materials, both living and dead, as well as dissolved colored compounds in the water column. During the melting of snow and ice in spring, lake water can become turbid (cloudy) from silt transported into the lake. Lake water usually clears in late spring but then becomes more turbid with increased algal growth as the summer progresses. The easiest and most widely used measure of lake water clarity is the Secchi disk depth.

Marie Lake is a moderately deep lake compared to many lakes in Alberta. Water clarity was measured four times in summer 2009, and compared to shallower lakes, Marie Lake is relatively clear with an average Secchi depth of 3.4 m in 2009. That summer, light penetrated to an average 13% of the total lake depth (**Table 1**), allowing algal growth in the top 6.8 m of the water column. The seasonal maximum Secchi depth of 4.0 m was recorded on 27 July and 30 August, and the seasonal minimum Secchi depth of 2.5 m was observed on 7 July.

Water Chemistry

Based on lake water characteristics, Marie Lake is considered mesotrophic (see *A Brief Introduction to Limnology* at the end of this report). In 2009, Marie Lake had moderate concentrations of total phosphorus (average TP = 19.6 µg/L), total nitrogen (average TN = 623.8 µg/L), and algal biomass (average chlorophyll *a* concentration = 3.26 µg/L). Total nitrogen concentrations varied from a seasonal minimum of 0.565 mg/L on 27 July to a seasonal maximum of 0.689 mg/L on 30 August (**Figure 5**). Total phosphorous declined over the summer from 32 µg/L on 7 June to a seasonal minimum of 13 µg/L on 17 August. Chlorophyll *a* (a measure of algal biomass) concentrations were relatively constant over the open water period, and varied from a maximum of 5.82 µg/L on 7 June to a minimum of 2.22 µg/L on 27 July. Marie Lake does not appear to be impacted by nutrient loading associated with human use and land use activities in the watershed.

Marie Lake is well protected from acidification; its pH of 8.5 is well above that of pure water (i.e., pH 7). Ion concentrations have remained virtually unchanged in Marie Lake since data were collected in 1980 (**Table 1**), although carbonate and bicarbonate were not measured prior to 2007. The stable ion concentrations suggest the hydrology of Marie Lake has remained in equilibrium during the sampling years, and stable sulfate concentrations indicate that atmospheric deposition of acidic petroleum-related pollutants does not currently influence the chemistry of runoff into Marie Lake.

The concentrations of various metals (as total recoverable concentrations) in Marie Lake were measured twice in 2009. All metal concentrations were below CCME guidelines for the Protection of Freshwater Aquatic Life (**Appendix 1**).




Figure 5. Total phosphorous, chlorophyll *a* (a measure of algal biomass), and total nitrogen concentrations for Marie Lake during the summer of 2009.

Table 1. Mean water chemistry and Secchi depth values from Marie Lake, Alberta, for summer 2009 compared to previous years.

Parameter	1980	1981	2002	2003	2004	2007	2008	2009
TP ($\mu\text{g L}^{-1}$)	-	15	13	12	19	15	17.4	19.8
TDP ($\mu\text{g L}^{-1}$)	-	8	5	4	3	5.5	6.6	11.3
Chl a ($\mu\text{g L}^{-1}$)	6.5	4.6	2.1	4	4.8	2.6	2.7	3.3
Secchi depth (m)	2.5	3.0	4.6	5.75	3.7	3.4	3.8	3.4
TN ($\mu\text{g L}^{-1}$)	-	-	517	495	550	495	552	624
NO ₂₊₃ ($\mu\text{g L}^{-1}$)	<1	-	1.8	4.8	5.2	<5	8	6.3
NH ₄ ⁺ ($\mu\text{g L}^{-1}$)	-	<22	9.3	6.5	9.0	9	15.2	16.3
Dissolved organic C (mg L^{-1})	-	-	-	-	-	10.8	11.4	10.4
Ca (mg L^{-1})	30	-	35	34	33	34.2	32.9	31.5
Mg (mg L^{-1})	12	-	12	14	12	12.2	12.2	11.7
Na (mg L^{-1})	6	-	6	6	6.6	6.9	6.4	6.5
K (mg L^{-1})	2	-	2	2	2	2	1.9	1.7
SO ₄ ²⁻ (mg L^{-1})	< 3	-	0.69	<3	<3	<3	3.3	3.7
Cl ⁻ (mg L^{-1})	< 1	-	0.56	0.47	0.6	0.8	0.6	0.83
CO ₃ (mg L^{-1})	-	-	-	-	-	5	4	9
HCO ₃ (mg L^{-1})	-	-	-	-	-	171	173	165
Conductivity ($\mu\text{S cm}^{-1}$)	-	-	-	-	-	267	266	263
pH	-	-	-	-	8.44	8.4	8.4	8.5
TDS (mg L^{-1})	-	-	-	-	-	146.5	144.3	143
Total Alkalinity ($\text{mg L}^{-1} \text{CaCO}_3$)	135	-	147	152	171	148	146	146

Note: TP = total phosphorous, TDP = total dissolved phosphorous, TN = total Kjeldahl nitrogen, Chla = chlorophyll *a*, NO₂₊₃ = nitrate+nitrite, NH₄ = ammonium, Ca = calcium, Mg = magnesium, Na = sodium, K = potassium, SO₄ = sulphate, Cl = chloride, CO₃ = carbonate, HCO₃ = bicarbonate, Cond = Specific conductivity, TDS = Total dissolved solids.

References

- Mitchell, P. and E. Prepas. 1990. Atlas of Alberta Lakes. University of Alberta Press.
- Nürnberg, G.K. 1996. Trophic state of clear and colored, soft and hardwater lakes with special consideration of nutrients, anoxia, phytoplankton and fish. *Lake and Reservoir Management* 12(4):432-447.
- Strong, W.L. and K.R. Leggat. 1981. Ecoregions of Alberta. Alberta Energy and Natural Resources. 64 pp.
- Vollenweider, R.A., and J. Kerekes, J. 1982. Eutrophication of Waters. Monitoring, Assessment and Control. Organization for Economic Co-Operation and Development (OECD), Paris. 156 pp.
- Welch, E.B. 1980. Ecological Effects of Waste Water. Cambridge University Press.

Appendix 1

Mean concentrations of metals in Marie Lake, measured on 27 July and 30 August 2009, compared to average values for 2007 and 2008. Values shown for each year are the averages for each year. The CCME heavy metal Guidelines for the Protection of Freshwater Aquatic Life are presented for reference.

Metals (total)	2007	2008	2009	Guidelines
ALUMINUM µg/L	9.1	4.2	4.1	100 ^a
ANTIMONY µg/L	0.012	0.024	0.024	6 ^e
ARSENIC µg/L	0.51	0.67	0.68	5
BARIUM µg/L	33.3	32.1	31.1	1000 ^e
BERYLLIUM µg/L	<0.003	<0.003	<0.003	100 ^{d,f}
BISMUTH µg/L	<0.001	0.0013	0.0072	
BORON µg/L	17.7	23.0	24.3	5000 ^{e,f}
CADMIUM µg/L	<0.002	0.0029	0.0029	0.085 ^b
CHROMIUM µg/L	0.08	0.139	0.107	
COBALT µg/L	<0.001	0.0073	0.0075	1000 ^f
COPPER µg/L	0.13	<0.05	0.17	4 ^c
IRON µg/L	39.9	3.28	7.9	300
LEAD µg/L	0.021	0.0674	0.0395	7 ^c
LITHIUM µg/L	4.37	7.25	5.67	2500 ^g
MANGANESE µg/L	21.9	9.07	4.71	200 ^g
MOLYBDENUM µg/L	0.154	0.172	0.193	73 ^d
NICKEL µg/L	<0.005	0.086	0.058	150 ^c
SELENIUM µg/L	0.06	<0.1	0.158	1
STRONTIUM µg/L	90.8	91.0	90.5	
SILVER µg/L	<0.0005	<0.0005	<0.0005	
THALLIUM µg/L	<0.0003	<0.003	0.0015	0.8
THORIUM µg/L	0.007	<0.003	0.0009	
TIN µg/L	0.03	0.0315	<0.03	
TITANIUM µg/L	1.38	1.03	1.09	
URANIUM µg/L	0.08	0.0662	0.0667	100 ^e
VANADIUM µg/L	0.14	0.112	0.159	100 ^{f,g}
ZINC µg/L	0.64	0.175	0.162	30

Values represent means of total recoverable metal concentrations.

* one sample collected 27 June 2007.

^a Based on pH ≥ 6.5; calcium ion concentration [Ca⁺²] ≥ 4 mg/L; and dissolved organic carbon concentration [DOC] ≥ 2 mg/L.

^b Based on water Hardness of 300 mg/L (as CaCO₃).

^c Based on water Hardness > 180 mg/L (as CaCO₃).

^d CCME interim value.

^e Based of Canadian Drinking Water Quality guideline values.

^f Based of CCME Guidelines for Agricultural Use (Livestock Watering).

^g Based of CCME Guidelines for Agricultural Use (Irrigation).

A Brief Introduction to Limnology

Indicators of water quality

The goal of **Lakewatch** is to collect water samples necessary to determine the water quality of lakes. Though not all encompassing, the variables measured in **Lakewatch** are sensitive to human activities in watersheds that may cause impacts to water quality. For example, nutrients such as phosphorus and nitrogen are important determinants of lake productivity. The concentrations of these nutrients in a lake are affected (typically elevated) by land use changes such as increased crop production or livestock grazing. Elevated nutrient concentrations can cause increases in undesirable algae blooms resulting in low dissolved oxygen concentrations, degraded fish habitat and production of noxious odors. Large increases in nutrients over time may also indicate sewage inputs, which in turn, may result in other human health concerns such as harmful bacteria or protozoans (e.g. *Cryptosporidium*).

Temperature and mixing

Water temperature in a lake dictates the behavior of many chemical parameters responsible for water quality (Figure 6). Heat is transferred to a lake at its surface and slowly moves downward depending on water circulation in the lake. Lakes with a large surface area or a small volume tend to have greater mixing due to wind. In deeper lakes, circulation is not strong enough to move warm water to depths typically greater than 4 or 5 m and as a result cooler denser water remains at the bottom of the lake. As the difference in temperature between warm surface and cold deeper water increases, two distinct layers are formed. Limnologists call these layers of water the **epilimnion** at the surface and the **hypolimnion** at the bottom. A transition layer known as the metalimnion, which contains the effective wall separating top and bottom waters called a thermocline, separates the layers. A thermocline typically occurs when water temperature changes by more than one degree within one-meter depth. The hypolimnion and epilimnion do not mix, nor do elements such as oxygen supplied at the surface move downward into the hypolimnion. In the fall, surface waters begin to cool and eventually reach the same temperature as hypolimnetic water. At this point, the water mixes from top to bottom in what is called a **turnover** event. Surface water cools further as ice forms and again a thermocline develops this time with 4° C water at the bottom and 0° C water on the top.




Figure 6: Difference in the circulation of the water column depending on thermal stratification.

In spring, another turnover event occurs when surface waters warm to 4° C. Lakes with this mixing pattern of two stratification periods and two turnover events are called **dimictic** lakes. In shallower lakes, the water column may mix from top to bottom most of the ice-free season with occasional stratification during periods of calm warm conditions. Lakes that mix frequently are termed **polymictic** lakes. In our cold climate, many shallow lakes are **cold monomictic** meaning a thermocline develops every winter, there is one turnover event in spring but the remainder of the ice-free season the lake is polymictic.

Dissolved Oxygen

Oxygen enters a lake at the lake surface and throughout the water column when produced by photosynthesizing plants, including algae, in the lake. Oxygen is consumed within the lake by respiration of living organisms and decomposition of organic material in the lake sediments. In lakes that stratify (see temperature above), oxygen that dissolves into the lake at the surface cannot mix downward into the

hypolimnion. At the same time, oxygen is depleted in the hypolimnion by decomposition. The result is that the hypolimnion of a lake can become **anoxic**, meaning it contains little or no dissolved oxygen. When a lake is frozen, the entire water column can become anoxic because the surface is sealed off from the atmosphere. Winter anoxic conditions can result in a fish-kill, which is particularly common during harsh winters with extended ice-cover. Alberta Surface Water Quality Guidelines suggest dissolved oxygen concentrations (in the epilimnion) must not decline below 5 mg/L and should not average less than 6.5 mg/L over a seven-day period. However, the guidelines also require that dissolved oxygen concentrations remain above 9.5 mg/L in areas where early life stages of aquatic biota, particularly fish, are present.

General Water Chemistry

Water in lakes always contains substances that have been transported by rain and snow or have entered the lake in groundwater and inflow streams. These substances may be dissolved in the water or suspended as particles. Some of these substances are familiar minerals, such as sodium and chloride, which when combined form table salt, but when dissolved in water separate into the two electrically charged components called **ions**. Most dissolved substances in water are in ionic forms and are held in solution due to the polar nature of the water molecule. **Hydrophobic** (water-fearing) compounds such as oils contain little or no ionic character, are non-polar and for this reason do not readily dissolve in water. Although hydrophobic compounds do not readily dissolve, they can still be transported to lakes by flowing water. Within individual lakes, ion concentrations vary from year to year depending on the amount and mineral content of the water entering the lake. This mineral content can be influenced by the amount of precipitation and other climate variables as well as human activities such as fertilizer and road salt application.

Phosphorus and Nitrogen

Phosphorus and nitrogen are important nutrients limiting the growth of algae in Alberta lakes. While nitrogen usually limits agricultural plants, phosphorus is usually in shortest supply in lakes. Even a slight increase of phosphorus in a lake can, given the right conditions, promote algal blooms causing the water to turn green in the summer and impair recreational uses. When pollution originating from livestock manure and human sewage enters lakes, not only are the concentrations of phosphorus and nitrogen increased but nitrogen can become a limiting nutrient, which is thought to cause blooms of toxic algae belonging to the cyanobacteria. Not all cyanobacteria are toxic, however, the blooms can form decomposing mats that smell and impair dissolved oxygen concentrations in the lake.

Chlorophyll-a

Chlorophyll-*a* is a photosynthetic pigment that green plants, including algae, possess enabling them to convert the sun's energy to living material. Chlorophyll-*a* can be easily extracted from algae in the laboratory. Consequently, chlorophyll-*a* is a good estimate of the amount of algae in the water. Larger aquatic plants, known as macrophytes, rather than algae, dominate some highly productive lakes. In these lakes, chlorophyll-*a* and nutrient values taken from water samples do not include productivity from large aquatic plants. As a result, lakes like Chestermere, which are dominated by macrophytes, can exist at a lower trophic state than if macrophyte biomass was included. Unfortunately, the productivity and nutrient cycling contributions of macrophytes are difficult to sample accurately and are therefore not typically included in trophic state indices.

Secchi Disk Depth

Lakes that are clear are more attractive for recreation, whereas those that are turbid or murky are considered by lake users to have poor water quality. Secchi disk depth is the oldest, simplest, and quickest quantitative measure of water clarity. A Secchi disk is a black and white disk that is lowered down through the water column until it can no longer be seen. Secchi disk depth is the midpoint between the depth at which it disappears when lowered and reappears when it is pulled up again. The Secchi disk depth in lakes with high algal biomass will generally be shallow. However, algae do not only affect Secchi disk depth. High

concentrations of suspended sediments, particularly fine clays or glacial till, are common in plains or mountain reservoirs of Alberta. Mountain reservoirs may have exceedingly shallow Secchi disk depths despite low algal growth and nutrient concentrations.

The euphotic zone, calculated as twice the Secchi disk depth, is the portion of the water column that has sufficient light for aquatic plants to grow. Murky waters, with shallow Secchi depths, can prevent aquatic plants from growing on the lake bottom. Aquatic plants are important because they ensure clear lake water by reducing shoreline erosion and stabilizing lake bottom sediments. Many lakes in Alberta are shallow and have bottom sediments with high concentrations of nutrients. Without aquatic plants, water quality may decline in these lakes due to murky, sediment-laden water and excessive algal blooms. Maintaining aquatic plants in certain areas of a lake is often essential for ensuring good water clarity and a healthy lake as many organisms, like aquatic invertebrates and fish, depend on aquatic plants for food and shelter.

Trophic state

Trophic state is a classification system for lakes that depends on fertility and is a useful index for rating and comparing lakes. From low to high nutrient and algal biomass (as chlorophyll-*a*) concentrations, the trophic states are: **oligotrophic**, **mesotrophic**, **eutrophic** and **hypereutrophic**. The nutrient and algal biomass concentrations that define these categories are shown in table 2 and a graph of Alberta lakes compared by trophic state can be found on the ALMS website. A majority of lakes in Alberta are meso- to eutrophic because they naturally contain high nutrient concentrations due to our deep fertile soils. Thus, lakes in Alberta are susceptible to human impacts because they are already nutrient-rich; any further nutrient increases can bring about undesirable conditions illustrated in Figure. 7.




Figure 7: Suggested changes in various lake characteristics with eutrophication. From “Ecological Effects of Wastewater”, 1980.

Table 2: Trophic status based on lake water characteristics

Trophic state	Total Phosphorus (µg/L)	Total Nitrogen (µg/L)	Chlorophyll a (µg/L)	Secchi Depth (m)
Oligotrophic	< 10	< 350	< 3.5	> 4
Mesotrophic	10 - 30	350 - 650	3.5 - 9	4 - 2
Eutrophic	30 - 100	650 - 1200	9 - 25	2 - 1
Hypereutrophic	> 100	> 1200	> 25	< 1

Note: These values are from a detailed study of global lakes reported in Nurnberg 1996. Alberta Environment uses slightly different values for TP and CHL based on those of the OECD reported by Vollenweider (1982). The AENV and OECD cutoffs for TP are 10, 35 and 100; for CHL are 3, 8 and 25. AENV does not have TN or Secchi depth criteria. The corresponding OECD exists for Secchi depth and the cutoffs are 6, 3 and 1.5 m.