



Preston McEachern
9637 – 81 Ave.
Edmonton, AB, T6C 0X6
(780) 427-1197
prestonm@telusplanet.net

Lakewatch

Sylvan Lake



*The Alberta Lake Management Society
Volunteer Lake Monitoring Report*

And you really live by the river? What a jolly life!"

"By it and with it and on it and in it," said the Rat. "It's brother and sister to me. What it hasn't got is not worth having, and what it doesn't know is not worth knowing." Kenneth Grahame The Wind in the Willows

"The world's supply of fresh water is running out. Already one person in five has no access to safe drinking water." BBC World Water Crisis Homepage

A note from the Lakewatch Coordinator

Preston McEachern

Lakewatch has several important objectives, one of which is to document and interpret water quality in Alberta Lakes. Equally important are the objectives of educating lake users about their aquatic environment; enhancing public involvement in lake management; and facilitating a link between aquatic scientists and lake users. The Lakewatch Reports are designed to summarize basic lake data in understandable terms for a lay audience, and are not meant to be a complete synopsis of information about specific lakes. Substantial additional information is generally available on the lakes that have participated in Lakewatch and readers requiring more information are encouraged to seek these sources.

The 2002 Lakewatch Report has undergone a substantial change in format from previous years. I am no longer the author as much as an editor including text and figures from others who have done an excellent job describing lakes throughout Alberta. I have attempted to give due credit to these outstanding people and apologize for blatant plagiarism where it occurs. As editor, feel free to castigate me for errors. I have included easily accessible information that is likely to have been updated in recent years and readers are encouraged to help update these reports by sending new information to me.

I would like to thank all people who share my love for aquatic environments and particularly those who have helped in the Lakewatch program. These people prove that ecological apathy can be overcome and give us hope that water will not be the limiting factor in the health of our planet.

Acknowledgements

The Lakewatch program is made possible through the dedication of its volunteers and Alberta Environment employees. Mike Bilyk, John Willis, Doreen LeClair and Dave Trew from Alberta Environment were instrumental in funding, training people and organizing with Lakewatch data. Comments on this report by Dave Trew were appreciated. Alberta Lake Management Society members and the board of directors helped in many facets of water collection and management. Sophie Lewin and Lucille Kowalchuk were summer field coordinators and were an excellent addition to the program. Without the dedication of these people and the interest of cottage owners, Lakewatch would not have occurred.

Sylvan Lake

Sylvan Lake is a large (42.8 km²), moderately deep (maximum depth 18.3 m) lake just West of the city of Red Deer. The lake was first named “Snake Lake” from the Indian name *Kinabik*, which referred to the numerous garter snakes in the area. The name was officially changed to Sylvan Lake in 1903. “Sylvan” is from the Latin *sylvanus*, which means “of a forest”. Most of the surrounding land was originally forested with trembling aspen. However, approximately 90% of the watershed has been converted to agriculture.

Sylvan Lake was first settled in 1899 and, because of its rare natural beauty, it soon started to become a summer resort by 1904. Since then, the shore of Sylvan Lake has become intensively developed with four summer villages, the town of Sylvan Lake and six subdivisions.

Two provincial parks also occupy the lakeshore: Jarvis Bay and Sylvan Lake. Majestic vertical sandstone banks rise to a height of up to 20 m along the northeast shore. In general, the lake’s shoreline is mostly sand or a mixture of rock and gravel.

Rooted aquatic plants occur in patches in sheltered areas around the lake and grow densely in the northwest end. The most common emergent species are bulrush (*Scirpus* sp.) and common cattail (*Typha latifolia*). Submergent macrophytes, which grow to a depth of 3.5 m, include pondweeds (*Potamogeton* spp.), water buttercup (*Ranunculus circinata*), Canada water weed (*Elodea canadensis*) and the macroalgae stonewort (*Chara* sp.). There are at least seven species of fish in Sylvan Lake: northern pike, yellow perch, walleye, burbot, spottail shiners, brook stickleback, and fathead minnows.




Fig. 1: Bathymetry of Sylvan Lake (Mitchell and Prepas 1990).




Fig. 2: Water levels in Sylvan Lake for the period of 1960 to 2001.

Water Levels

The drainage basin of Sylvan Lake likely contributes little to water levels since it is small compared to the area of the lake (only 2.5 times the area of the lake). This is further supported by the intermittent nature of the inflowing streams and the presence of numerous submerged springs. Evaporation of Sylvan Lake's large surface area seems to be the primary outlet for water since very little water flows out of Sylvan Lake: the outlet stream flowed during part of only three years between 1955 and 1976.

The water level of Sylvan Lake has been recorded regularly since 1955 and periodically as far back as 1918. The elevation of Sylvan Lake's surface is quite stable; from 1960 to 2001 the range was only about 1 meter (Fig 3). The maximum lake level throughout recorded history was 937.2 m in 1955. The minimum lake level occurred in 1939 when it reached 936.0 m. In 2001, water levels were close to the 1960-2001 average at 936.7. Water levels in Sylvan Lake seem to have been decreasing over the last 10 years. However, since the early 1960s, water levels have actually increased slightly.




Fig. 3 & 4: Temperature and dissolved oxygen (DO) profiles for Sylvan Lake for the summer 2001.

Results

Water Temperature and Dissolved Oxygen

Water temperature and dissolved oxygen profiles in the water column can provide information on water quality and fish habitat. Please refer to the end of this report for descriptions of technical terms.

At the end of June 2001, a weak thermocline developed at about 1.5 meters depth. Over the next two months, thermal stratification destabilized: in July, the thermocline deepened to about 10 meters and did not form in August. Also, water temperatures were 3 to 5 degrees warmer in the two last months, as compared to June. Oxygen concentrations in June were steady with depth except for the bottom two meters where they declined drastically. In July and August, dissolved oxygen concentrations decreased at depths greater than 10 meters, reflecting increased microbial metabolism in the warmer waters of mid/late summer.

Water clarity and Secchi Depth

Water clarity is influenced by suspended materials, both living and dead, as well as some coloured dissolved compounds in the water column. During the melting of snow and ice in spring, lake water can become cloudy from silt transported into the lake. Lake water usually clears in late




Fig. 5 & 6: Total phosphorus, chlorophyll *a* and Kjeldahl nitrogen for Sylvan Lake, summer 2001.

spring but then becomes more turbid with increased algal biomass as the summer progresses. The easiest and most widely used measure of lake water clarity is the Secchi disk depth.

Sylvan Lake's water was clear during the summer of 2001. Secchi disk depth was 3 m in the beginning of June and increased progressively to 5 m by the end of August. This increase in water clarity over the summer is surprising given that algal biomass also increased during this time. Factors other than algal biomass were responsible for this change of water clarity in Sylvan Lake.

Water chemistry

Because Sylvan Lake had moderate nutrient concentrations and algal biomass compared to lakes throughout Canada, it is considered mesotrophic (see details on trophic status classification at end of this report). In the Alberta context, Sylvan Lake is a little below average in these characteristics. In 2001, total phosphorus concentrations were fairly stable and hovered around 20 µg/L for most of the summer but then increased substantially (i.e.,

Table 1: Mean values from summer 2001 samples compared to values from those reported in the Atlas of Alberta Lakes.

Parameter	1984	1986	2001
Total phosphorus (µg/L)	20	21	23
TDP (µg/L)	N/A	N/A	8
Chlorophyll <i>a</i> (µg/L)	3.8	3.7	9
Secchi disk depth (m)	5.0	4.7	
Total nitrogen (µg/L)	-	-	836
TDN (µg/L)	-	-	489
NO ₂₊₃ (µg/L)	-	-	2
NH ₄ (µg/L)	-	-	9
Ca (mg/L)	-	18	17
Mg (mg/L)	-	37	36
Na (mg/L)	-	64	60
K (mg/L)	-	7	7
SO ₄ (mg/L)	-	16	13
Cl (mg/L)	-	<1	3
Alkalinity (mg/L CaCO ₃)	-	325	318
Conductivity (µS/cm)	-	597	572
pH	-	8.8-9.0	9
Colour (mg/L Pt)	-	-	3
TSS (mg/L)	-	-	2

Note. TDP = total dissolved phosphorus, TDN = total dissolved nitrogen, NO₂₊₃ = nitrate+nitrite, NH₄⁺ = ammonium, Ca = calcium, Mg = magnesium, Na = sodium, K = potassium, SO₄ = sulfate, Cl = chloride, TSS = total suspended solids.

about 2 times) at the end of August. Chlorophyll *a*, a measure of algal biomass, mirrored changes in total phosphorus concentrations (Fig. 5). This pattern of increased phosphorus and chlorophyll *a* concentrations in Sylvan Lake towards the end of summer is typical of most lakes in Alberta. Sylvan Lake was moderately concentrated in nitrogen with TKN ranging from 427 to 1487 µg/L and averaging 834 µg/L. In general, the water quality of Sylvan Lake was relatively good and the water was clear.

Sylvan Lake is well buffered: its pH of 9 is well above that of pure water (=7). Its dominant ions are bicarbonate, sodium, and magnesium. The relatively high sodium and magnesium concentrations indicate substantial groundwater inflow. Ion concentrations were very similar to historic values from the Atlas of Alberta Lakes. In general, water quality did not change much over the last 15 years.

A brief introduction to Limnology

Indicators of water quality

Water samples are collected in Lakewatch to determine the water quality of lakes. Though not all encompassing, the variables collected in Lakewatch are sensitive to human activities in watersheds that can cause degraded water quality. For example, nutrients such as phosphorus and nitrogen are important determinants of lake productivity. The concentrations of these nutrients in a lake are impacted (typically elevated) by land use changes such as increased crop production or livestock grazing. Elevated nutrient concentrations can cause increases in undesirable algae blooms resulting in low dissolved oxygen concentrations, degraded habitat for fish and noxious smells. A large increase in nutrients over time may also indicate sewage inputs which in turn may result in other human health concerns associated with bacteria or the protozoan *Cryptosporidium*.

Temperature and mixing

Water temperature in a lake dictates the behavior of many chemical parameters responsible for water quality. Heat is transferred to a lake at its surface and slowly moves downward depending on water circulation in the lake. Lakes with a large surface area or a small volume tend to have greater mixing due to wind. In deeper lakes, circulation is not strong enough to move warm water to depths typically greater than 4 or 5 m and as a result cooler denser water remains at the bottom of the lake. As the difference in temperature

between warm surface and cold deeper water increases, two distinct layers are formed. Limnologists call these layers of water the **epilimnion** at the surface and the **hypolimnion** at the bottom. The layers are separated by a transition layer known as the **metalimnion** which contains the effective wall separating top and bottom waters called a **thermocline**. A thermocline typically occurs when water temperature changes by more than one degree within one meter depth. The hypolimnion and epilimnion do not mix, nor do elements such as oxygen supplied at the surface move downward into the hypolimnion. In the fall, surface waters begin to cool and eventually reach the same temperature as hypolimnetic water. At this point the water mixes from top to bottom in what is called a **turnover** event. Surface water cools further as ice forms and again a thermocline develops this time with 4° C water at the bottom and 0° C water on the top.

In spring another turnover event occurs when surface waters warm to 4° C. Lakes with this mixing pattern of two stratification periods and two turnover events are called **dimictic** lakes. In shallower lakes, the water column may mix from top to bottom most of the ice-free season with occasional stratification during periods of calm warm conditions. Lakes that mix frequently are termed **polymictic** lakes. In our cold climate, many shallow lakes are **cold monomictic** meaning a thermocline develops every winter, there is one turnover event in spring but the remainder of the ice free season the lake is polymictic.

Dissolved Oxygen

Oxygen enters a lake at the lake surface and throughout the water column when produced by photosynthesizing plants, including algae, in the lake. Oxygen is consumed within the lake by




Fig. 7: Difference in the circulation of the water column depending on thermal stratification.

respiration of living organisms and decomposition of organic material in the lake sediments. In lakes that stratify (see temperature above), oxygen that dissolves into the lake at the surface cannot mix downward into the hypolimnion. At the same time oxygen is depleted in the hypolimnion by decomposition. The result is that the hypolimnion of a lake can become **anoxic**, meaning it contains little or no dissolved oxygen. When a lake is frozen, the entire water column can become anoxic because the surface is sealed off from the atmosphere. Winter anoxic conditions can result in a fish-kill which is particularly common during harsh winters with extended ice-cover. Alberta Surface Water Quality Guidelines suggest dissolved oxygen concentrations (in the epilimnion) must not decline below 5 mg/L and should not average less than 6.5 mg/L over a seven-day period. However, the guidelines also require that dissolved oxygen concentrations remain above 9.5 mg/L in areas where early life stages of aquatic biota, particularly fish, are present.

General Water Chemistry

Water in lakes always contains substances that have been transported by rain and snow or have entered the lake in groundwater and inflow streams. These substances may be dissolved in the water or suspended as particles. Some of these substances are familiar minerals, such as sodium and chloride, which when combined form table salt, but when dissolved in water separate into the two electrically charged components called **ions**. Most dissolved substances in water are in ionic forms and are held in solution due to the polar nature of the water molecule. **Hydrophobic** (water-fearing) compounds such as oils contain little or no ionic character, are non-polar and for this reason do not readily dissolve in water. Although hydrophobic compounds do not readily dissolve, they can still be transported to lakes by flowing water. Within individual lakes, ion concentrations vary from year to year depending on the amount and mineral content of the water entering the lake. This mineral content can be influenced by the amount of precipitation and other climate variables as well as human activities such as fertilizer and road salt application.

Phosphorus and Nitrogen

Phosphorus and nitrogen are important nutrients limiting the growth of algae in Alberta lakes. While nitrogen usually limits agricultural plants, phosphorus is usually in shortest supply in lakes. Even a slight increase of phosphorus in a lake can, given the right conditions, promote algal blooms causing the water to turn green in the summer and impair recreational uses. When pollution originating from livestock manure and human sewage enters lakes not only are the concentrations of phosphorus and nitrogen increased but nitrogen can become a limiting nutrient which is thought to cause blooms of toxic algae belonging to the cyanobacteria. Not all cyanobacteria are toxic, however, the blooms can form decomposing mats that smell and impair dissolved oxygen concentrations in the lake.

Chlorophyll a

Chlorophyll *a* is a photosynthetic pigment that green plants, including algae, possess enabling them to convert the sun's energy to living material. Chlorophyll *a* can be easily extracted from algae in the laboratory. Consequently, chlorophyll *a* is a good estimate of the amount of algae in the water. Some highly productive lakes are dominated by larger aquatic plants, known as macrophytes, rather than suspended algae. In these lakes, chlorophyll *a* and nutrient values taken from water samples do not include productivity from large aquatic plants. As a result, lakes like Chestermere which are dominated by macrophytes can be at a lower trophic state than if macrophyte biomass was included. Unfortunately, the productivity and nutrient cycling contributions of macrophytes are difficult to sample accurately and are therefore not typically included in trophic state indices.

Secchi Disk Depth

Lakes that are clear are more attractive for recreation, whereas those that are turbid or murky are considered by lake users to have poor water quality. Secchi disk depth is the oldest, simplest, and

quickest quantitative measure of water clarity. A Secchi disk is a black and white disk that is lowered down through the water column until it can no longer be seen. Secchi disk depth is the midpoint between the depth at which it disappears when lowered and reappears when it is pulled up again. The Secchi disk depth in lakes with high algal biomass will generally be shallow. However, Secchi disk depth is not only affected by algae. High concentrations of suspended sediments, particularly fine clays or glacial till, are common in plains or mountain reservoirs of Alberta. Mountain reservoirs may have exceedingly shallow Secchi disk depths despite low algal growth and nutrient concentrations.

The euphotic zone, calculated as twice the Secchi disk depth, is the portion of the water column that has sufficient light for aquatic plants to grow. Murky waters, with shallow Secchi depths, can prevent aquatic plants from growing on the lake bottom. Aquatic plants are important because they ensure clear lake water by reducing shoreline erosion and stabilizing lake bottom sediments. Many lakes in Alberta are shallow and have bottom sediments with high concentrations of nutrients. Without aquatic plants, water quality may decline in these lakes due to murky, sediment-laden water and excessive algal blooms. Maintaining aquatic plants in certain areas of a lake is often essential for ensuring good water clarity and a healthy lake as many organisms, like aquatic invertebrates and fish, depend on aquatic plants for food and shelter.

Trophic state

Trophic state is a classification system for lakes that depends on fertility and is a useful index for rating and comparing lakes. From low to high nutrient and algal biomass (as chlorophyll) concentrations, the trophic states are: **oligotrophic**, **mesotrophic**, **eutrophic** and **hypereutrophic**. The nutrient and algal biomass concentrations that define these categories are shown in the following table, a figure of Alberta lakes compared by trophic state can be found on the ALMS website. A majority of lakes in Alberta are meso- to eutrophic because they naturally contain high nutrient concentrations due to our deep fertile soils. Thus, lakes in Alberta are susceptible to human impacts because they are already nutrient-rich; any further nutrient increases can bring about undesirable conditions illustrated in Fig 8.




Fig. 8: Suggested changes in various lake characteristics with eutrophication. From “Ecological Effects of Wastewater”, 1980

Trophic status based on lake water characteristics.

Trophic state	Total Phosphorus (µg/L)	Total Nitrogen (µg/L)	Chlorophyll a (µg/L)	Secchi Depth (m)
Oligotrophic	< 10	< 350	< 3.5	> 4
Mesotrophic	10 - 30	350 - 650	3.5 - 9	4 - 2
Eutrophic	30 - 100	650 - 1200	9 - 25	2 - 1
Hypereutrophic	> 100	> 1200	> 25	< 1

Note: These values are from a detailed study of global lakes reported in Nurnberg 1996. Alberta Environment uses slightly different values for TP and CHL based on those of the OECD reported by Vollenweider (1982). The AENV and OECD cutoffs for TP are 10, 35 and 100; for CHL are 3, 8 and 25. AENV does not have TN or Secchi depth criteria. The corresponding OECD exists for Secchi depth and the cutoffs are 6, 3 and 1.5 m.