



*THE ALBERTA LAKE MANAGEMENT SOCIETY
VOLUNTEER LAKE MONITORING PROGRAM*

2015 Hanmore Lake Report

LAKEWATCH IS MADE POSSIBLE WITH SUPPORT FROM:



Ce projet a été réalisé avec l'appui financier de :
This project was undertaken with the financial support of:



Environnement
Canada

Environment
Canada





Alberta Lake Management Society's LakeWatch Program

LakeWatch has several important objectives, one of which is to collect and interpret water quality data on Alberta Lakes. Equally important is educating lake users about their aquatic environment, encouraging public involvement in lake management, and facilitating cooperation and partnerships between government, industry, the scientific community and lake users. LakeWatch Reports are designed to summarize basic lake data in understandable terms for a lay audience and are not meant to be a complete synopsis of information about specific lakes. Additional information is available for many lakes that have been included in LakeWatch and readers requiring more information are encouraged to seek those sources.

ALMS would like to thank all who express interest in Alberta's aquatic environments and particularly those who have participated in the LakeWatch program. These people prove that ecological apathy can be overcome and give us hope that our water resources will not be the limiting factor in the health of our environment.

This report has been published before the completion of the data validation process.

Acknowledgements

The LakeWatch program is made possible through the dedication of its volunteers. We would like to thank Dean & Carol Milne for their assistance with sampling Hanmore Lake in 2015. We would also like to thank Laticia McDonald, Ageleky Bouzetos, and Mohamad Youssef who were summer technicians with ALMS in 2015. Executive Director Bradley Peter was instrumental in planning and organizing the field program. Mike Bilyk was involved in the training aspects of the program. Lisa Reinbolt was responsible for data management. This report was prepared by Bradley Peter and Alicia Kennedy. The Beaver River Watershed Alliance (BRWA), the Lakeland Industry and Community Association (LICA), the Alberta Environmental Monitoring Evaluation and Reporting Agency (AEMERA), and Environment Canada, were major sponsors of the program.

HANMORE LAKE:

Hanmore Lake is a moderately small lake located in the North Saskatchewan Watershed. It is 20 km north of the town of Smoky Lake and 120 km northwest of Edmonton. Two public campgrounds are located on Hanmore Lake: Hanmore Lake West is located on the south side of the west end of the lake, and Hanmore Lake East is located on the far east end of the lake in the south bay. Pioneer Bible Camp is located on the south shore. There is only one agricultural plot bordering the lake on the northwest side of the lake. The name was first recorded in 1920 under Little Whitefish Lake because of its large whitefish as well as its close proximity to Whitefish Lake¹.



Figure 1 – Looking East down the length of Hanmore Lake. Photo by Jackson Woren.

The lake is long and narrow, with a length of 4.2 km and a maximum width of 1 km. Drop-offs in Hanmore Lake are fairly steep and the maximum depth is around 23 m in three holes just west of the center of the lake. There are six ephemeral streams that flow into Hanmore Lake during large rainstorm events and spring runoff.

Hanmore Lake is primarily used for recreational purposes for activities such as swimming, boating, canoeing, kayaking, paddle boarding, and other water sports. Sportfishing is popular at Hanmore Lake and species include northern pike, whitefish, and yellow perch². It is not known what the macrophyte composition is at Hanmore Lake but it is known that the lake perimeter includes species such as the common cattail (*Typha latifolia*) as well as bulrushes (*Scirpus spp.*). Unidentified submerged species are most dense in the west bay, as well as in the shallow north east bay.

WATER QUANTITY:

There are many factors influencing water quantity. Some of these factors include the size of the lakes drainage basin, precipitation, evaporation, water consumption, ground water influences, and the efficiency of the outlet channel structure at removing water from the lake. Requests for water quantity monitoring should go through Environment and Sustainable Resource Developments Monitoring and Science division.

¹ Merrily K. Aubrey. 2006. Concise Place Names of Alberta. University of Calgary Press Available at: <http://www.albertasource.ca/placenames>

² Alberta Fishing Guide. 2014. Available at: <http://albertafishingguide.com/>

The water level at Hanmore Lake has not changed dramatically between 1969 and 2013². In 1993 the water level was at its lowest at 668.62 meters above sea level (m asl) and peak water levels were observed at 670.08 m asl in 1974 (Figure 2). It is believed that the increase in water levels in 1974 was attributed to above normal snowfalls and heavier than normal spring rains. It is not known how long the water residency time is at Hanmore Lake. More recent validated water level data was not available.




Figure 2 - Water levels in elevation as meters above sea levels (m asl) from 1969 to 2013. Data obtained from Alberta Environment and Sustainable Resource Development.

WATER CLARITY AND SECCHI DEPTH:

Water clarity is influenced by suspended materials, both living and dead, as well as dissolved colored compounds in the water column. During the melting of snow and ice in spring, lake water can become turbid (cloudy) from silt transported into the lake. Lake water usually clears in late spring but then becomes more turbid with increased algal growth as the summer progresses. The easiest and most widely used measure of lake water clarity is the Secchi disk depth.

Hanmore Lake had an average Secchi disk depth of 3.22 m in 2015 (Table 1). This average is only slightly less than that observed in 2014 and is comprised of a maximum observed clarity of 4.25 m on June 15th and a minimum observed clarity of 2.00 m on August 19th (Figure 3). Water clarity decreased from June 15th to August 19th, though it showed a recovery on September 21st.

² University of Alberta. 2005. Atlas of Alberta Lakes; Hubbles Lake, Lake Basin Characteristics. University of Alberta Press. Available at: <http://sunsite.ualberta.ca/Projects/Alberta-Lakes/>

Water clarity may have been impaired by a lake whiting event, in which extremely small phytoplankton cause precipitation of calcium onto their surface, giving the lake a glacial green appearance (technician observation). Though chlorophyll-*a* concentrations were low, the field technician did observe cyanobacteria on the lake's surface which may have also contributed to reduced water clarity.




Figure 3 – Secchi depth values measured five times over the course of the summer at Hanmore Lake in 2015.

WATER TEMPERATURE AND DISSOLVED OXYGEN:

Water temperature and dissolved oxygen profiles in the water column can provide information on water quality and fish habitat. The depth of the thermocline is important in determining the depth to which dissolved oxygen from the surface can be mixed. Please refer to the end of this report for descriptions of technical terms.

As a deep lake, Hanmore shows strong thermal stratification throughout the summer (Figure 4a). This results in warm surface waters and cool bottom waters. Thermal stratification remained strong until September 12th when a weakening of the thermocline was observed. It is unknown whether Hanmore Lake mixes during the fall. At the surface, a maximum temperature of 21.88 °C was observed on July 13th. Below the thermocline, water temperatures regularly proceeded to approximately 6 °C. Hanmore's thermal stratification has important implications for its dissolved oxygen concentrations.




Figure 4 – a) Temperature (°C) and b) dissolved oxygen (mg/L) profiles for Hanmore Lake measured five times over the course of the summer of 2015. Drifting on September 12th caused the field technician to miss the ‘deep spot’.

The surface of Hanmore Lake remained well oxygenated throughout the summer. Surface concentrations ranged between 8.73 mg/L on September 12th and 9.65 mg/L on June 15th (Figure 4b). These concentrations fall well above the Canadian Council for Ministers of the Environment guidelines for the Protection of Aquatic Life of 6.5 mg/L. At the thermocline, a spike in dissolved oxygen was observed on two of the five sampling trips – this spike may be indicative of a metalimnetic bloom which is likely responsible for the lake whiting event. Below the thermocline, oxygen concentrations regularly proceeded toward anoxia. This is likely a combination of decomposition occurring at the sediments and isolation from well-oxygenated surface waters. Anoxic conditions at the sediment may promote the release of phosphorus from the sediment into overlying waters.

WATER CHEMISTRY:

ALMS measures a suite of water chemistry parameters. Phosphorus, nitrogen, and chlorophyll-a are important because they are indicators of eutrophication, or excess nutrients, which can lead to harmful algal/cyanobacteria blooms. One direct measure of harmful cyanobacteria blooms are Microcystins, a common group of toxins produced by cyanobacteria. See Table 1 for a complete list of parameters.

Average total phosphorus concentration (TP) measured 12 µg/L in 2015 (Table 1). This value falls into the mesotrophic, or moderately productive, classification and is very similar to that measured in 2014 (Table 1). Throughout the summer, TP fluctuated from a maximum of 14 µg/L on both June 15th and August 9th, and a minimum of 10 µg/L on August 19th (Figure 5). Phosphorus input into Hanmore Lake should be limited in order to maintain Hanmore Lake's water quality.

Average chlorophyll-*a* concentration measured 2.9 µg/L in 2015 (Table 1). This value falls into the oligotrophic, or low productivity, classification. This value is very similar to that measured in 2014 (Table 1) and was comprised of a maximum of 3.6 mg/L on July 13th and 2.4 mg/L on September 21st (Figure 5). The field technician did observe cyanobacteria, specifically *Aphanizomenon* spp., though this was not detected in significant quantities.

Finally, total Kjeldahl nitrogen (TKN) measured 0.69 in 2015 (Table 1). This value falls into the eutrophic, or productive, classification. Many lakes in Alberta have elevated levels of nitrogen – however, total phosphorus is the primary nutrient driving phytoplankton blooms.




Figure 5 – Total Phosphorus (TP), Total Kjeldahl Nitrogen (TKN), and Chlorophyll-*a* concentration measured five times over the course of the summer at Hanmore Lake.

Average pH measured 8.65 in 2015 – this value is well above neutral. Hanmore Lake is well buffered against changes to pH due to its alkalinity (212 mg/L CaCO₃) and bicarbonate (240 mg/L HCO₃) concentrations. Hanmore Lake has moderate conductivity (382 uS/cm) with magnesium (27 mg/L), sodium (22 mg/L), and calcium (21 mg/L) as dominant contributing ions.

Metals were collected twice at Hanmore Lake and all concentrations fell within their respective guidelines (Table 2).

MICROCYSTIN:

Microcystins are toxins produced by cyanobacteria (blue-green algae) which, when ingested, can cause severe liver damage. Microcystins are produced by many species of cyanobacteria which are common to Alberta's Lakes, and are thought to be the one of the most common cyanobacteria toxins. In Alberta, recreational guidelines for microcystin are set at 20 µg/L.

In 2015, all tests for microcystin fell below the laboratory's minimum detection limits (0.1 µg/L; table 1).

INVASIVE SPECIES:

Quagga and Zebra mussels are invasive species which, if introduced to our lakes, will have significant negative ecological, economical, and recreational impacts. ALMS collects water samples which are analyzed for mussel veligers (juveniles) and monitors substrates for adult mussels. In order to prevent the spread of invasive mussels, always clean, drain, and dry your boat between lakes. To report mussel sightings or mussel-fouled boats, call the confidential Alberta hotline at 1-855-336-BOAT.

In 2015, no invasive zebra or quagga mussels were detected in Hanmore Lake.

Table 1 - Average Secchi disk depth and water chemistry values for Hanmore Lake. Previous years averages are provided for comparison.

Parameter	1986	2014	2015
TP (µg/L)	16	13	12
TDP (µg/L)	/	7	6
Chl- <i>a</i> (µg/L)	/	2.4	2.9
Secchi depth (m)	/	3.89	3.22
TKN (mg/L)	/	0.92	0.69
NO ₂ and NO ₃ (µg/L)	0.025	20	2.5
NH ₃ (µg/L)	/	28	25
DOC (mg/L)	/	9.4	9.4
Ca (mg/L)	31	20	21
Mg (mg/L)	19	25	27
Na (mg/L)	11	23	22
K (mg/L)	6.1	7.8	8.1
SO ₄ ²⁻ (mg/L)	2.5	2.3	2.1
Cl ⁻ (mg/L)	0.5	0.9	1
CO ₃ (mg/L)	7.2	13	10
HCO ₃ (mg/L)	202.35	249	240
pH	8.7	8.67	8.65
Conductivity (µS/cm)	333	374	382
Hardness (mg/L)	155	155	162
TDS (mg/L)	179.96	210	208
Microcystin (µg/L)	/		<0.1
Total Alkalinity (mg/L CaCO ₃)	178	204	212

Note: TP = total phosphorous, TDP = total dissolved phosphorous, Chl-*a* = chlorophyll-*a*, TKN = total Kjeldahl nitrogen. NO₂₊₃ = nitrate+nitrite, NH₃ = ammonia, Ca = calcium, Mg = magnesium, Na = sodium, K = potassium, SO₄ = sulphate, Cl = chloride, CO₃ = carbonate, HCO₃ = bicarbonate, TDS = total dissolved solids. A forward slash (/) indicates an absence of data.

Table 2 - Average concentrations of metals measured in Hanmore Lake on August 9th and September 21st. Values shown for 2015 are an average of those dates. The CCME heavy metal Guidelines for the Protection of Freshwater Aquatic Life (unless otherwise indicated) are presented for reference.

Metals (Total Recoverable)	2014	2015	Guidelines
Aluminum µg/L	8.75	10.5	100 ^a
Antimony µg/L	0.037	0.026	6 ^e
Arsenic µg/L	0.994	0.989	5
Barium µg/L	44.5	45.4	1000 ^e
Beryllium µg/L	0.004	0.004	100 ^{d,f}
Bismuth µg/L	0.0005	0.002	/
Boron µg/L	86.2	92.4	5000 ^{ef}
Cadmium µg/L	0.0025	0.001	0.085 ^b
Chromium µg/L	0.3	0.085	/
Cobalt µg/L	0.007	0.014	1000 ^f
Copper µg/L	0.205	1.73	4 ^c
Iron µg/L	29.45	22.4	300
Lead µg/L	0.080	0.032	7 ^c
Lithium µg/L	30.2	33.7	2500 ^g
Manganese µg/L	9.88	10.71	200 ^g
Molybdenum µg/L	0.132	0.2305	73 ^d
Nickel µg/L	0.004	0.004	150 ^c
Selenium µg/L	0.03	0.03	1
Silver µg/L	0.001	0.001	0.1
Strontium µg/L	166	173	/
Thallium µg/L	0.00045	0.00045	0.8
Thorium µg/L	0.003825	0.006125	/
Tin µg/L	0.0085	0.06275	/
Titanium µg/L	0.315	0.425	/
Uranium µg/L	0.322	0.325	100 ^e
Vanadium µg/L	0.06	0.06	100 ^{f,g}
Zinc µg/L	0.35	0.6	30

Values represent means of total recoverable metal concentrations.

^a Based on pH ≥ 6.5; calcium ion concentrations [Ca⁺²] ≥ 4 mg/L; and dissolved organic carbon concentration [DOC] ≥ 2 mg/L.

^b Based on water Hardness of 300 mg/L (as CaCO₃)

^c Based on water hardness > 180mg/L (as CaCO₃)

^d CCME interim value.

^e Based on Canadian Drinking Water Quality guideline values.

^f Based on CCME Guidelines for Agricultural use (Livestock Watering).

^g Based on CCME Guidelines for Agricultural Use (Irrigation).

A forward slash (/) indicates an absence of data or guidelines.

A BRIEF INTRODUCTION TO LIMNOLOGY

INDICATORS OF WATER QUALITY:

Water samples are collected in LakeWatch to determine the chemical characteristics that characterize general water quality. Though not all encompassing, the variables collected in LakeWatch are sensitive to human activities in watersheds that can cause degraded water quality. For example, nutrients such as phosphorus and nitrogen are important determinants of lake productivity. The concentrations of these nutrients in a lake are impacted (typically elevated) by land use changes such as increased crop production or livestock grazing. Elevated nutrient concentrations can cause increases in undesirable algae blooms resulting in low dissolved oxygen concentrations, degraded habitat for fish and noxious smells. A large increase in nutrients over time may also indicate sewage inputs which in turn may result in other human health concerns associated with bacteria or the protozoan *Cryptosporidium*.

TEMPERATURE AND MIXING:

Water temperature in a lake dictates the behavior of many chemical parameters responsible for water quality. Heat is transferred to a lake at its surface and slowly moves downward depending on water circulation in the lake. Lakes with a large surface area or a small volume tend to have greater mixing due to wind. In deeper lakes, circulation is not strong enough to move warm water to depths typically greater than 4 or 5 m and as a result cooler denser water remains at the bottom of the lake. As the difference in temperature between warm surface and cold deeper water increases, two distinct layers are formed. Limnologists call these layers of water the **epilimnion** at the surface and the **hypolimnion** at the bottom. The layers are separated by a transition layer known as the **metalimnion** which contains the effective wall separating top and bottom waters called a **thermocline**. A thermocline typically occurs when water temperature changes by more than one degree within one meter depth. The hypolimnion and epilimnion do not mix, nor do elements such as oxygen supplied at the surface move downward into the hypolimnion. In the fall, surface waters begin to cool and eventually reach the same temperature as hypolimnetic water. At this point the water mixes from top to bottom in what is often called a **turnover** event. Surface water cools further as ice forms and again a thermocline develops this time with 4° C water at the bottom and near 0° C water on the top.




Figure A: Difference in the circulation of the water column depending on thermal stratification.

In spring another turnover event occurs when surface waters warm to 4° C. Lakes with this mixing pattern of two stratification periods and two turnover events are called **dimictic** lakes. In shallower lakes, the water column may mix from top to bottom most of the ice-free season with occasional stratification during periods of calm warm conditions. Lakes that mix frequently are **termed polymictic** lakes. In our cold climate, many shallow lakes are **cold monomictic meaning a thermocline develops every winter, there is one turnover event in spring but the remainder of the ice free season the lake is polymictic.**

DISSOLVED OXYGEN:

Oxygen enters a lake at the lake surface and throughout the water column when produced by photosynthesizing plants, including algae, in the lake. Oxygen is consumed within the lake by respiration of living organisms and decomposition of organic material in the lake sediments. In lakes that stratify (see temperature above), oxygen that dissolves into the lake at the surface cannot mix downward into the hypolimnion. At the same time oxygen is depleted in the hypolimnion by decomposition. The result is that the hypolimnion of a lake can become **anoxic**, meaning it contains little or no dissolved oxygen. When a lake is frozen, the entire water column can become anoxic because the surface is sealed off from the atmosphere. Winter anoxic conditions can result in a fish-kill which is particularly common during harsh winters with extended ice-cover. Alberta Surface Water Quality Guidelines suggest dissolved oxygen concentrations (in the epilimnion) must not decline below 5 mg•L⁻¹ and should not average less than 6.5 mg•L⁻¹ over a seven-day period. However, the guidelines also require that dissolved oxygen concentrations remain above 9.5 mg•L⁻¹ in areas where early life stages of aquatic biota, particularly fish, are present.

GENERAL WATER CHEMISTRY:

Water in lakes always contains substances that have been transported by rain and snow or have entered the lake in groundwater and inflow streams. These substances may be dissolved in the water or suspended as particles. Some of these substances are familiar minerals, such as sodium and chloride, which when combined form table salt, but when dissolved in water separate into the two electrically charged components called **ions**. Most dissolved substances in water are in ionic forms and are held in solution due to the polar nature of the water molecule. **Hydrophobic** (water-fearing) compounds such as oils contain little or no ionic character, are non-polar and for this reason do not readily dissolve in water. Although hydrophobic compounds do not readily dissolve, they can still be transported to lakes by flowing water. Within individual lakes, ion concentrations vary from year to year depending on the amount and mineral content of the water entering the lake. This mineral content can be influenced by the amount of precipitation and other climate variables as well as human activities such as fertilizer and road salt application.

PHOSPHORUS AND NITROGEN:

Phosphorus and nitrogen are important nutrients limiting the growth of algae in Alberta lakes. While nitrogen usually limits agricultural plants, phosphorus is usually in shortest supply in lakes. Even a slight increase of phosphorus in a lake can, given the right conditions, promote algal blooms causing the water to turn green in the summer and impair recreational uses. When

pollution originating from livestock manure and human sewage enters lakes not only are the concentrations of phosphorus and nitrogen increased but nitrogen can become a limiting nutrient which is thought to cause blooms of toxic algae belonging to the cyanobacteria. Not all cyanobacteria are toxic, however, the blooms can form decomposing mats that smell and impair dissolved oxygen concentrations in the lake.

CHLOROPHYLL-A:

Chlorophyll *a* is a photosynthetic pigment that green plants, including algae, possess enabling them to convert the sun's energy to living material. Chlorophyll *a* can be easily extracted from algae in the laboratory. Consequently, chlorophyll *a* is a good estimate of the amount of algae in the water. Some highly productive lakes are dominated by larger aquatic plants rather than suspended algae. In these lakes, chlorophyll *a* and nutrient values taken from water samples do not include productivity from large aquatic plants. The result, in lakes like Chestermere which are dominated by larger plants known as macrophytes, can be a lower trophic state than if macrophyte biomass was included. Unfortunately, the productivity and nutrient cycling contributions of macrophytes are difficult to sample accurately and are therefore not typically included in trophic state indices.

SECCHI DISK TRANSPARENCY :

Lakes that are clear are more attractive for recreation, whereas those that are turbid or murky are considered by lake users to have poor water quality. A measure of the transparency or clarity of the water is performed with a Secchi disk with an alternating black and white pattern. To measure the clarity of the water, the Secchi disk is lowered down into the water column and the depth where the disk disappears is recorded. The Secchi depth in lakes with a lot of algal growth will be small while the Secchi depth in lakes with little algal growth can be very deep. However, low Secchi depths are not caused by algal growth alone. High concentrations of suspended sediments, particularly fine clays or glacial till, are common in plains or mountain reservoirs of Alberta. Mountain reservoirs may have exceedingly low Secchi depths despite low algal growth and nutrient concentrations.

The euphotic zone or the maximum depth that light can penetrate into the water column for actively growing plants is calculated as twice the Secchi depth. Murky waters, with shallow Secchi depths, can prevent aquatic plants from growing on the lake bottom. Conversely, aquatic plants can ensure lakes have clear water by reducing shoreline erosion and stabilizing lake bottom sediments. In Alberta, many lakes are shallow and bottom sediments contain high concentrations of nutrients. Without aquatic plants, water quality may decline in these lakes due to murky, sediment laden water and excessive algal blooms. Maintaining aquatic plants in certain areas of a lake is often essential for ensuring good water clarity and a healthy lake as many organisms, like aquatic invertebrates and insects, depend on aquatic plants for food and shelter.

TROPHIC STATE:

Trophic state is classification of lakes into four categories of fertility and is a useful index for rating and comparing lakes. From low to high nutrient and algal biomass (as chlorophyll) concentrations, the trophic states are; **oligotrophic, mesotrophic, eutrophic** and **hypereutrophic** (Table 2).

A majority of lakes in Alberta contain naturally high levels of chlorophyll *a* (8 to 25 µg/L) due to our deep fertile soils. These lakes are usually considered fertile and are termed eutrophic. The nutrient and algal biomass concentrations that define these categories are shown in the following table, a figure of Alberta lakes compared by trophic state can be found on the ALMS website.




Figure B: Suggested changes in various lake characteristics with eutrophication. From “Ecological Effects of Wastewater”, 1980.

Table A - Trophic status classification based on lake water characteristics.

Trophic state	Total Phosphorus (µg•L ⁻¹)	Total Nitrogen (µg•L ⁻¹)	Chlorophyll a (µg•L ⁻¹)	Secchi Depth (m)
Oligotrophic	< 10	< 350	< 3.5	> 4
Mesotrophic	10 – 30	350 - 650	3.5 - 9	4 - 2
Eutrophic	30 – 100	650 - 1200	9 - 25	2 - 1
Hypereutrophic	> 100	> 1200	> 25	< 1