



*The Alberta Lake Management Society
Volunteer Lake Monitoring Program*

Little Beaver Lake Tributary Report 2018

Lakewatch is made possible
with support from:

This project was undertaken with the financial support of:
Ce projet a été réalisé avec l'appui financier de :



Environment and
Climate Change Canada

Environnement et
Changement climatique Canada



Environment
and Parks



Cargill



Beaver River Watershed Alliance





ALBERTA LAKE MANAGEMENT SOCIETY'S LAKEWATCH PROGRAM

LakeWatch has several important objectives, one of which is to collect and interpret water quality data on Alberta Lakes. Equally important is educating lake users about their aquatic environment, encouraging public involvement in lake management, and facilitating cooperation and partnerships between government, industry, the scientific community and lake users. LakeWatch Reports are designed to summarize basic lake data in understandable terms for a lay audience and are not meant to be a complete synopsis of information about specific lakes. Additional information is available for many lakes that have been included in LakeWatch and readers requiring more information are encouraged to seek those sources.

ALMS would like to thank all who express interest in Alberta's aquatic environments and particularly those who have participated in the LakeWatch program. These people prove that ecological apathy can be overcome and give us hope that our water resources will not be the limiting factor in the health of our environment.



ACKNOWLEDGEMENTS

The LakeWatch program is made possible through the dedication of its volunteers. We would like to extend a special thanks to Tony Cable and Doug Jensen of the Friends of Little Beaver Lake Society for helping to initiate this project at Little Beaver Lake. We would also like to acknowledge the Little Beaver Lake Stewardship Society for their financial contributions to this project. Field sampling and report writing was conducted by Bradley Peter and Laura Redmond.

INTRODUCTION

Little Beaver Lake is a quiet, scenic lake 35 km south of Camrose and 107 km south of Edmonton in the Battle River Watershed. This shallow lake is approximately 3.5 km long and 500 m wide, and is surrounded by forested rolling hills and agricultural development. The county subdivision of Little Beaver Lake Estates lies on its west shore, and the Village of Ferintosh lies on its east shore.

The lake surface area is 1.66 km² and the lake watershed is approximately 63 km². Little Beaver Lake's lake:watershed ratio is 1:38 (Figure 1). The northern tributary was sampled in 2017 and flows from a similarly sized lake to Little Beaver Lake through a culvert undercutting Highway 21 just south of the railroad tracks. The southern tributary flows from the south, under Township Road 435A, into Little Beaver Lake just west of the outflow.

The purpose of this study was to determine the loading (amount) of key nutrients entering Little Beaver Lake through the southern tributary from April 20 to May 9, 2018.



The south tributary of Little Beaver Lake on April 19, 2018.

HOW WERE DATA COLLECTED?

The downstream side of the southern culvert was sampled three times (Table 1) in the spring of 2018 for information related to flow and water chemistry.

Flow: The width of the culvert and depth of the stream were used to calculate the cross-sectional area of flow. The velocity of the stream was measured using a Gurley Price meter connected to a Wading Rod. This device uses rotating cups held below the surface of the stream to measure velocity. The area of flow and stream velocity were used to calculate discharge in m³/s. Discharge was assumed constant until the subsequent sampling event.

Water Chemistry: Water samples were collected from below the surface and tested for various parameters, including: total phosphorus (TP), total dissolved phosphorus (TDP), nitrate and nitrite

(NO₃+NO₂), total Kjeldahl nitrogen (TKN), and ammonia. These parameters include various forms of phosphorus and nitrogen, nutrients which are important for growth in Little Beaver Lake. Water chemistry was assumed constant until the subsequent sampling event. Concentrations of these parameters were combined with discharge to calculate the loadings (mass) of nutrients passing through the culvert. Daily loadings were multiplied by the number of days between sampling events to determine total mass loading of each nutrient into the lake over the sampling period.

Table 1. Dates of culvert sampling and number of days between sampling visits.

Date	Days Between Samples
20-Apr	5
25-Apr	8
3-May	6
9-May	N/A*

*No flow on this date. Assumed end of flow.

RESULTS AND DISCUSSION:

Stream flow was highly variable from April 20-May 3. High velocity flow, which entirely filled the culvert, may have reduced the accuracy of flow measurements. The cross-sectional area of flow ranged from 0.91-0.92 m², while the velocity ranged from 0.08-0.93 m/s. These measurements resulted in a minimum discharge of 0.075 m³/s on May 3 and a maximum discharge of 0.85 m³/s on April 20 (Table 2). On May 9, water remained in the culvert though no flow was detected.

Table 2. Discharge (m³/s) and concentration of various nutrients (mg/L) observed on each sampling date in the southern tributary of Little Beaver Lake.

Date	Discharge (m ³ /s)	TP (mg/L)	TDP (mg/L)	NO ₃ +NO ₂ (mg/L)	TKN (mg/L)	Ammonia (mg/L)
20-Apr	0.85	0.80	0.65	1.9	2.9	0.51
25-Apr	0.28	0.52	0.46	2.1	2.4	0.42
3-May	0.075	0.51	0.46	0.0042	2.6	0.037
9-May	-	-	-	-	-	-

The total volume and mass loadings for the sampling season were calculated (Table 3). In sum, 6.0×10^8 L passed through the culvert between April 20-May 9, transporting 414 kg of total phosphorus, of which 345 kg were dissolved, 1106 kg of nitrate + nitrite, and 1631 kg of total Kjeldahl nitrogen including 270 kg of ammonia.

Table 3. Volume of water (L) and mass of nutrients (kg) passing through the southern tributary culvert at Little Beaver Lake.

Date	Volume (L)	TP (kg)	TDP (kg)	NO ₃ +NO ₂ (kg)	TKN (kg)	Ammonia (kg)
April 20-April 25	3.7×10^8	293.4	238.4	696.7	1063.4	187.0
April 25-May 3	1.9×10^8	101.2	89.5	408.6	467	81.7
May 3-May 9	3.9×10^7	19.8	17.8	0.2	100.7	1.4
Total April 20-May 9	6.0×10^8	414.3	345.7	1105.5	1631.1	270.2

Given that sampling events occurred at different intervals, the average concentration of nutrients in the stream over the period of sampling can be calculated as a flow weighted mean (Table 4).

Table 4. Flow weighted mean concentrations (mg/L) of each nutrient for the north tributary sampled in 2017 compared to the south tributary sampled in 2018.

Tributary	Year	TP (mg/L)	TDP (mg/L)	NO ₃ +NO ₂ (mg/L)	TKN (mg/L)	Ammonia (mg/L)
North	2017	0.723	0.633	0.233	2.58	0.131
South	2018	0.690	0.576	1.84	2.72	0.450

Concentrations of nutrients in the stream were high, often exceeding concentrations observed in the lake during summer sampling in previous years. For example, the 2016 average total phosphorus concentration in Little Beaver Lake was 0.168 mg/L compared to an average concentration of 0.69 mg/L observed in the southern tributary. Compared to the north tributary sampling from 2017, flow weighted concentrations of all nutrients in the south tributary were higher. Specifically, nitrate + nitrite levels were almost 7 x higher in the south tributary. High nutrient concentrations are likely attributed to the melting of winter snow stores which carry nutrients from the landscape. High water levels were observed at the south tributary, with the culvert being completely submerged on the first two sampling dates. Wetlands on the south side of Little Beaver Lake may help reduce nutrient concentrations before tributary water enters the lake, and the close proximity of the outflow may reduce the impact that this drainage has on the lake's summer conditions. Beneficial management practices applied to the landscape in Little Beaver Lake's watershed may result in improvements to Little Beaver Lake's water quality.



A large population of amphipods observed at the southern culvert in 2018.

FUTURE DIRECTIONS:

Further studies are needed to gain a more holistic view of the Little Beaver Lake watershed and inflows. This study has been combined with the sampling of the northern tributary in 2017, as well as a full lake survey in the summer of 2018. A greater number of sampling visits, including sampling during summer storm events, would increase the precision of mass loading measurements. Moreover, sampling across seasons would help provide clarity regarding the impact of seasonal variability. Future research may involve the preparation of a phosphorus budget including all tributaries leading to and from Little Beaver Lake.




Figure 1: Little Beaver Lake watershed boundary. Northern Tributary sampled in 2017 indicated by blue star. Southern Tributary sampled in 2018 indicated by yellow star.