



*THE ALBERTA LAKE MANAGEMENT SOCIETY
VOLUNTEER LAKE MONITORING PROGRAM*

2014 Lacombe Lake Report

LAKEWATCH IS MADE POSSIBLE WITH SUPPORT FROM:



Ce projet a été réalisé avec l'appui financier de :
This project was undertaken with the financial support of:



Environnement
Canada

Environment
Canada



Alberta Lake Management Society's LakeWatch Program

LakeWatch has several important objectives, one of which is to collect and interpret water quality data on Alberta Lakes. Equally important is educating lake users about their aquatic environment, encouraging public involvement in lake management, and facilitating cooperation and partnerships between government, industry, the scientific community and lake users. LakeWatch Reports are designed to summarize basic lake data in understandable terms for a lay audience and are not meant to be a complete synopsis of information about specific lakes. Additional information is available for many lakes that have been included in LakeWatch and readers requiring more information are encouraged to seek those sources.

ALMS would like to thank all who express interest in Alberta's aquatic environments and particularly those who have participated in the LakeWatch program. These people prove that ecological apathy can be overcome and give us hope that our water resources will not be the limiting factor in the health of our environment.

Acknowledgements

The LakeWatch program is made possible through the dedication of its volunteers. We would like to thank all the members of the Lacombe Lake Watershed Stewardship Society who assisted with sampling Lacombe Lake in 2014, including: Anto Davis, Elaine Atkinson Jones, Cliff Soper, and Ed Zaparniuk. We would also like to thank Jackson Woren, Brittany Kereliuk, and Kara MacAulay who were summer technicians with ALMS in 2014. Program Coordinator Bradley Peter was instrumental in planning and organizing the field program. Mike Bilyk was involved in the training aspects of the program. Lisa Reinbolt was responsible for data management. This report was prepared by Jackson Woren and Bradley Peter. Alberta Environment, the Beaver River Watershed Alliance (BRWA), the Lakeland Industry and Community Association (LICA), and Environment Canada, were major sponsors of the program.

LACOMBE LAKE:

Lacombe Lake is a pothole lake found in Lacombe County in central Alberta. It is located 5 km north of the town of Blackfalds and 15 km north of Red Deer. There are no public campgrounds around the lake as most of the land is private farms and homesteads as well as public land and reserves. It is thought that the lake was once called Jackfish Lake due to the northern pike found in the lake, though in 1975 the name was changed to Lacombe Lake. The Lacombe Lake area is part of the Treaty 6 Nations and was an area where the Samson and Erminskin Cree Nations hunted and travelled. Permanent camps were traditionally located in wooded areas as well as along rivers, and a known trade existed just south of Gull Lake.



Figure 1 –Photo of Lacombe Lake. Photo by Brittany Kereliuk, 2014.

The lake is long and narrow, with a length of about 3 km, a maximum depth of 2.9 m, and a maximum width of about 500 m. Lacombe Lake has numerous bays and points which give it a distinct shape. It is not known to be a popular fishing destination but the lake is used for non-motorized recreational water sports as well as swimming. Lacombe Lake is found in the Aspen Parkland ecoregion of Alberta, much of which is now farmland with other foliage such as trembling aspen, oak, mixed tall shrubs, and intermittent fescue grasslands⁴.

Known sportfish species at Lacombe lake are the northern pike, though angling websites state that other species may include walleye, burbot, whitefish, rainbow trout, brown trout, and brook trout¹. Lacombe Lake has a large population of macrophytes, including yellow pond lily, Richardson's pondweed, stonewort, cattail, bulrushes, and bladderwort. Due to its small size, dense macrophytes, and limited recreational activity, waterfowl are known to frequent the lake. Known species include the mallard, common grebe, goldeneye, scaup, and ruddy duck¹. Larger vertebrates that are found around the lake are deer, muskrat, and beavers.

In the 1960s, the Prairie Farm Rehabilitation Association constructed a weir on Whelp Creek to control and direct the flow into the north end of Lacombe Lake during periods of high flow. In the years previous to 2008, residents noticed that the water quality of Lacombe Lake was starting to deteriorate, as well as excessive macrophytes growth across the lake. The diversion was stopped and Golder Associates Ltd. assessed the water quality of Lacombe Lake over the period of 4 years. Golder Associates Ltd. concluded that the diversion may have brought in excess nutrients into Lacombe Lake and recommendations included enhanced monitoring, finding the

¹ <http://www.hookandbullet.com/fishing-lacombe-lake-blackfalds-ab/> 2015

² Golder Associates Ltd. 2013. Lacombe Lake Water Quality Assessment - Alberta

³ Ecoregions of Canada. 1995. Available at: <http://ecozones.ca/English/region/156.html>

source of nutrients and bacteria, and improving water quality through best management practices. Best management practices includes watershed controls such as fertilizer restrictions, restoration of riparian vegetation along shorelines, and nutrient management planning.

WATER QUANTITY:

There are many factors influencing water quantity. Some of these factors include the size of the lakes drainage basin, precipitation, evaporation, water consumption, ground water influences, and the efficiency of the outlet channel structure at removing water from the lake. Requests for water quantity monitoring should go through Environment and Sustainable Resource Developments Monitoring and Science division.

No water quantity data has been collected at Lacombe Lake; therefore, the connections between surface water and groundwater are not yet understood.

WATER CLARITY AND SECCHI DEPTH:

Water clarity is influenced by suspended materials, both living and dead, as well as dissolved colored compounds in the water column. During the melting of snow and ice in spring, lake water can become turbid (cloudy) from silt transported into the lake. Lake water usually clears in late spring but then becomes more turbid with increased algal growth as the summer progresses. The easiest and most widely used measure of lake water clarity is the Secchi disk depth. Two times the Secchi disk depth equals the euphotic depth – the depth to which there is enough light for photosynthesis.

Throughout the sampling season, Secchi disk depth did not change dramatically at Lacombe Lake (Figure 3). On June 6th, Secchi disk depth measured a seasonal maximum of 2.2 m, while on September 7th, Secchi disk depth measured a seasonal minimum of 1.15 m. This decrease in water clarity correlated closely with increases in chlorophyll-*a* concentration, suggesting phytoplankton is the primary factor impeding water clarity at Lacombe Lake. On average, Secchi disk depth measured 1.53 m in 2014.




Figure 2 – Secchi disk depths recorded from the profile spot at Lacombe Lake, 2014.

WATER TEMPERATURE AND DISSOLVED OXYGEN:

Water temperature and dissolved oxygen profiles in the water column can provide information on water quality and fish habitat. The depth of the thermocline is important in determining the depth to which dissolved oxygen from the surface can be mixed. Please refer to the end of this report for descriptions of technical terms.

Surface water temperatures ranged from a minimum of 16.42 °C on September 7th to a maximum of 23.59 °C on August 14th (Figure 3a). Lacombe Lake remained isothermal on each sampling trip, with slight changes to water temperatures near the sediment. Due to the shallow depths at Lacombe Lake, thermal stratification may be observed on calm, extreme temperature days, though the lake likely mixes multiple times throughout the year. The presence/absence of thermal stratification has important implications for nutrient cycling and dissolved oxygen concentrations in Lacombe Lake.




Figure 3 – a) Temperature (°C) and b) dissolved oxygen (mg/L) profiles from Lacombe Lake, 2014.

Surface dissolved oxygen concentrations did not vary at Lacombe Lake during 2014 (Figure 3b). On all sampling dates, oxygen concentrations at the surface remained well above the Canadian Council for Ministers of the Environment (CCME) Guidelines for the Protection of Aquatic Life (PAL) of 6.5 mg/L. Surface concentrations measured a minimum of 8.92 mg/L on June 6th and a maximum of 9.63 mg/L on September 7th in 2014. Dissolved oxygen concentrations decreased rapidly as they approached the sediment. Anoxic conditions near the sediment are likely the result of decomposition – this may have important implications for the release of phosphorus from the lake sediments.

WATER CHEMISTRY:

*ALMS measures a suite of water chemistry parameters. Phosphorus, nitrogen, and chlorophyll-*a* are important because they are indicators of eutrophication, or excess nutrients, which can lead to harmful algal/cyanobacteria blooms. One direct measure of harmful cyanobacteria blooms are Microcystins, a common group of toxins produced by cyanobacteria. See Table 1 for a more complete list of parameters.*

Average Total Phosphorus (TP) measured 22.05 $\mu\text{g/L}$ in 2014 (Table 1). This value falls into the mesotrophic, or moderately productive, classification. Throughout the summer, TP ranged from a seasonal minimum of 18.6 $\mu\text{g/L}$ on September 7th to a seasonal maximum of 27.6 $\mu\text{g/L}$ on July 5th (Figure 4). Not enough data exists to establish nutrient trends at Lacombe Lake.

Chlorophyll-*a* concentration measured an average of 7.68 $\mu\text{g/L}$ in 2014 (Table 1). This value falls well into the mesotrophic classification. Throughout the summer, chlorophyll-*a* concentration ranged from a minimum of 4 $\mu\text{g/L}$ on June 6th to a maximum of 12 $\mu\text{g/L}$ on September 7th (Figure 4). More data is required to better understand the relationship between nutrients and chlorophyll-*a* concentration at Lacombe Lake.

Finally, total Kjeldahl nitrogen (TKN) measured an average of 1308 $\mu\text{g/L}$ in 2014 (Table 1). Unlike phosphorus and chlorophyll-*a*, average TKN falls into the hypereutrophic classification. Throughout the summer, TKN ranged from a minimum of 1250 $\mu\text{g/L}$ on July 5th and September 7th to a maximum of 1370 $\mu\text{g/L}$ on August 17th (Figure 4).




Figure 4 – Total phosphorus ($\mu\text{g/L}$), chlorophyll-*a* ($\mu\text{g/L}$), and total Kjeldahl nitrogen ($\mu\text{g/L}$) concentrations measured four times over the course of the summer of 2014.

Average pH measured 8.535 in 2014 – this value is well above neutral. Lacombe Lake has high alkalinity (227.83 mg/L CaCO₃) and bicarbonate (261 mg/L HCO₃) concentration which will help to buffer the lake against changes to pH (Table 2). Sodium (34.08 mg/L), magnesium (31.61 mg/L), and calcium (27.18 mg/L) are the dominant ions in Lacombe Lake, contributing to a conductivity of 505.67 µS/cm. More data needs to be collected to understand historical trends (Table 1).

Metals were collected twice at Lacombe Lake and all concentrations fell within their respective guidelines (Table 2).

MICROCYSTIN:

Microcystins are toxins produced by cyanobacteria (blue-green algae) which, when ingested, can cause severe liver damage. Microcystins are produced by many species of cyanobacteria which are common to Alberta's Lakes, and are thought to be the one of the most common cyanobacteria toxins. In Alberta, recreational guidelines for microcystin are set at 20 µg/L.

In 2014, concentrations of microcystin reached an observed maximum of 0.19 µg/L on September 7th (Table 1) – well below the Alberta recreational guidelines. It is unclear which cyanobacteria species were dominant throughout the open water season in 2014. Caution should always be taken when recreating in waters experiencing cyanobacteria blooms.

INVASIVE SPECIES:

Quagga and Zebra mussels are invasive species which, if introduced to our lakes, will have significant negative ecological, economical, and recreational impacts. ALMS collects water samples which are analyzed for mussel veligers (juveniles) and monitors substrates for adult mussels. In order to prevent the spread of invasive mussels, always clean, drain, and dry your boat between lakes. To report mussel sightings or mussel-fouled boats, call the confidential Alberta hotline at 1-855-336-BOAT.

In 2014, no zebra or quagga mussels were detected in Lacombe Lake.

Table 1 – Average Secchi disk depth and water chemistry values for Lacombe Lake. Limited data from previous years is provided for comparison.

Parameter	2008 ^a	2009 ^a	2010 ^a	2011 ^a	2012 ^a	2014
TP (µg/L)	*	*	*	45.67	16.4	22.05
TDP (µg/L)	\	\	\	\	\	5.12
Chlorophyll- <i>a</i> (µg/L)	\	\	\	\	\	7.68
Secchi depth (m)	\	\	\	\	\	1.54
TKN (µg/L)	1125	1866.67	1920	1600	1280	1308.333
NO ₂ and NO ₃ (µg/L)	20	35	13.2	4.5	3	5.28
NH ₃ (µg/L)	*	*	*	45.6	82	18.02
DOC (mg/L)	\	\	\	\	\	\
Ca (mg/L)	\	\	\	\	\	27.18
Mg (mg/L)	\	\	\	\	\	31.62
Na (mg/L)	\	\	\	\	\	34.08
K (mg/L)	\	\	\	\	\	12.63
SO ₄ ²⁻ (mg/L)	\	\	\	\	\	14.18
Cl ⁻ (mg/L)	\	\	\	\	\	21.45
CO ₃ (mg/L)	\	\	\	\	\	8.12
HCO ₃ (mg/L)	\	\	\	\	\	261
pH	\	\	\	\	\	8.54
Conductivity (µS/cm)	\	\	\	\	\	505.67
Hardness (mg/L)	\	\	\	\	\	198
TDS (mg/L)	\	\	\	\	\	278.17
Microcystin (µg/L)	\	\	\	\	\	0.15
Total Alkalinity (mg/L CaCO ₃)	\	\	\	\	\	227.83

Note: TP = total phosphorus, TDP = total dissolved phosphorus, Chl-*a* = chlorophyll-*a*, TKN = total Kjeldahl nitrogen. NO₂₊₃ = nitrate+nitrite, NH₃ = ammonia, Ca = calcium, Mg = magnesium, Na = sodium, K = potassium, SO₄ = sulphate, Cl = chloride, CO₃ = carbonate, HCO₃ = bicarbonate. A forward slash (/) indicates an absence of data.

^a Data from Golder Associates Ltd. 2013. Lacombe Lake Water Quality Assessment - Alberta

*Data excluded due to high detection limits.

Table 2 - Concentrations of metals measured in Lacombe Lake on June 6th and September 7th, 2014. The CCME heavy metal Guidelines for the Protection of Freshwater Aquatic Life (unless otherwise indicated) are presented for reference.

Metals (Total Recoverable)	2014	Guidelines
Aluminum µg/L	14	100 ^a
Antimony µg/L	0.0595	6 ^c
Arsenic µg/L	0.9115	5
Barium µg/L	62.25	1000 ^e
Beryllium µg/L	0.004	100 ^{d,f}
Bismuth µg/L	0.0005	/
Boron µg/L	45.75	5000 ^{ef}
Cadmium µg/L	0.0015	0.085 ^b
Chromium µg/L	0.175	/
Cobalt µg/L	0.033	1000 ^f
Copper µg/L	0.3975	4 ^c
Iron µg/L	17.7	300
Lead µg/L	0.01475	7 ^c
Lithium µg/L	19.8	2500 ^g
Manganese µg/L	48.1	200 ^g
Molybdenum µg/L	0.137	73 ^d
Nickel µg/L	0.042	150 ^c
Selenium µg/L	0.175	1
Silver µg/L	0.001	0.1
Strontium µg/L	199.5	/
Thallium µg/L	0.001575	0.8
Thorium µg/L	0.001975	/
Tin µg/L	0.00775	/
Titanium µg/L	0.865	/
Uranium µg/L	0.6785	100 ^e
Vanadium µg/L	0.185	100 ^{f,g}
Zinc µg/L	0.95	30

Values represent means of total recoverable metal concentrations.

^a Based on pH ≥ 6.5; calcium ion concentrations [Ca⁺²] ≥ 4 mg/L; and dissolved organic carbon concentration [DOC] ≥ 2 mg/L.

^b Based on water Hardness of 300 mg/L (as CaCO₃)

^c Based on water hardness > 180mg/L (as CaCO₃)

^d CCME interim value.

^e Based on Canadian Drinking Water Quality guideline values.

^f Based on CCME Guidelines for Agricultural use (Livestock Watering).

^g Based on CCME Guidelines for Agricultural Use (Irrigation).

A forward slash (/) indicates an absence of data or guidelines.

A BRIEF INTRODUCTION TO LIMNOLOGY

INDICATORS OF WATER QUALITY:

Water samples are collected in LakeWatch to determine the chemical characteristics that characterize general water quality. Though not all encompassing, the variables collected in LakeWatch are sensitive to human activities in watersheds that can cause degraded water quality. For example, nutrients such as phosphorus and nitrogen are important determinants of lake productivity. The concentrations of these nutrients in a lake are impacted (typically elevated) by land use changes such as increased crop production or livestock grazing. Elevated nutrient concentrations can cause increases in undesirable algae blooms resulting in low dissolved oxygen concentrations, degraded habitat for fish and noxious smells. A large increase in nutrients over time may also indicate sewage inputs which in turn may result in other human health concerns associated with bacteria or the protozoan *Cryptosporidium*.

TEMPERATURE AND MIXING:

Water temperature in a lake dictates the behavior of many chemical parameters responsible for water quality. Heat is transferred to a lake at its surface and slowly moves downward depending on water circulation in the lake. Lakes with a large surface area or a small volume tend to have greater mixing due to wind. In deeper lakes, circulation is not strong enough to move warm water to depths typically greater than 4 or 5 m and as a result cooler denser water remains at the bottom of the lake. As the difference in temperature between warm surface and cold deeper water increases, two distinct layers are formed. Limnologists call these layers of water the **epilimnion** at the surface and the **hypolimnion** at the bottom. The layers are separated by a transition layer known as the **metalimnion** which contains the effective wall separating top and bottom waters called a **thermocline**. A thermocline typically occurs when water temperature changes by more than one degree within one meter depth. The hypolimnion and epilimnion do not mix, nor do elements such as oxygen supplied at the surface move downward into the hypolimnion. In the fall, surface waters begin to cool and eventually reach the same temperature as hypolimnetic water. At this point the water mixes from top to bottom in what is often called a **turnover** event. Surface water cools further as ice forms and again a thermocline develops this time with 4° C water at the bottom and near 0° C water on the top.




Figure A: Difference in the circulation of the water column depending on thermal stratification.

In spring another turnover event occurs when surface waters warm to 4° C. Lakes with this mixing pattern of two stratification periods and two turnover events are called **dimictic** lakes. In shallower lakes, the water column may mix from top to bottom most of the ice-free season with occasional stratification during periods of calm warm conditions. Lakes that mix frequently are termed **polymictic** lakes. In our cold climate, many shallow lakes are **cold monomictic** meaning a thermocline develops every winter, there is one turnover event in spring but the remainder of the ice free season the lake is polymictic.

DISSOLVED OXYGEN:

Oxygen enters a lake at the lake surface and throughout the water column when produced by photosynthesizing plants, including algae, in the lake. Oxygen is consumed within the lake by respiration of living organisms and decomposition of organic material in the lake sediments. In lakes that stratify (see temperature above), oxygen that dissolves into the lake at the surface cannot mix downward into the hypolimnion. At the same time oxygen is depleted in the hypolimnion by decomposition. The result is that the hypolimnion of a lake can become **anoxic**, meaning it contains little or no dissolved oxygen. When a lake is frozen, the entire water column can become anoxic because the surface is sealed off from the atmosphere. Winter anoxic conditions can result in a fish-kill which is particularly common during harsh winters with extended ice-cover. Alberta Surface Water Quality Guidelines suggest dissolved oxygen concentrations (in the epilimnion) must not decline below 5 mg•L⁻¹ and should not average less than 6.5 mg•L⁻¹ over a seven-day period. However, the guidelines also require that dissolved oxygen concentrations remain above 9.5 mg•L⁻¹ in areas where early life stages of aquatic biota, particularly fish, are present.

GENERAL WATER CHEMISTRY:

Water in lakes always contains substances that have been transported by rain and snow or have entered the lake in groundwater and inflow streams. These substances may be dissolved in the water or suspended as particles. Some of these substances are familiar minerals, such as sodium and chloride, which when combined form table salt, but when dissolved in water separate into the two electrically charged components called **ions**. Most dissolved substances in water are in ionic forms and are held in solution due to the polar nature of the water molecule. **Hydrophobic** (water-fearing) compounds such as oils contain little or no ionic character, are non-polar and for this reason do not readily dissolve in water. Although hydrophobic compounds do not readily dissolve, they can still be transported to lakes by flowing water. Within individual lakes, ion concentrations vary from year to year depending on the amount and mineral content of the water entering the lake. This mineral content can be influenced by the amount of precipitation and other climate variables as well as human activities such as fertilizer and road salt application.

PHOSPHORUS AND NITROGEN:

Phosphorus and nitrogen are important nutrients limiting the growth of algae in Alberta lakes. While nitrogen usually limits agricultural plants, phosphorus is usually in shortest supply in lakes. Even a slight increase of phosphorus in a lake can, given the right conditions, promote algal blooms causing the water to turn green in the summer and impair recreational uses. When

pollution originating from livestock manure and human sewage enters lakes not only are the concentrations of phosphorus and nitrogen increased but nitrogen can become a limiting nutrient which is thought to cause blooms of toxic algae belonging to the cyanobacteria. Not all cyanobacteria are toxic, however, the blooms can form decomposing mats that smell and impair dissolved oxygen concentrations in the lake.

CHLOROPHYLL-A:

Chlorophyll *a* is a photosynthetic pigment that green plants, including algae, possess enabling them to convert the sun's energy to living material. Chlorophyll *a* can be easily extracted from algae in the laboratory. Consequently, chlorophyll *a* is a good estimate of the amount of algae in the water. Some highly productive lakes are dominated by larger aquatic plants rather than suspended algae. In these lakes, chlorophyll *a* and nutrient values taken from water samples do not include productivity from large aquatic plants. The result, in lakes like Chestermere which are dominated by larger plants known as macrophytes, can be a lower trophic state than if macrophyte biomass was included. Unfortunately, the productivity and nutrient cycling contributions of macrophytes are difficult to sample accurately and are therefore not typically included in trophic state indices.

SECCHI DISK TRANSPARENCY :

Lakes that are clear are more attractive for recreation, whereas those that are turbid or murky are considered by lake users to have poor water quality. A measure of the transparency or clarity of the water is performed with a Secchi disk with an alternating black and white pattern. To measure the clarity of the water, the Secchi disk is lowered down into the water column and the depth where the disk disappears is recorded. The Secchi depth in lakes with a lot of algal growth will be small while the Secchi depth in lakes with little algal growth can be very deep. However, low Secchi depths are not caused by algal growth alone. High concentrations of suspended sediments, particularly fine clays or glacial till, are common in plains or mountain reservoirs of Alberta. Mountain reservoirs may have exceedingly low Secchi depths despite low algal growth and nutrient concentrations.

The euphotic zone or the maximum depth that light can penetrate into the water column for actively growing plants is calculated as twice the Secchi depth. Murky waters, with shallow Secchi depths, can prevent aquatic plants from growing on the lake bottom. Conversely, aquatic plants can ensure lakes have clear water by reducing shoreline erosion and stabilizing lake bottom sediments. In Alberta, many lakes are shallow and bottom sediments contain high concentrations of nutrients. Without aquatic plants, water quality may decline in these lakes due to murky, sediment laden water and excessive algal blooms. Maintaining aquatic plants in certain areas of a lake is often essential for ensuring good water clarity and a healthy lake as many organisms, like aquatic invertebrates and insects, depend on aquatic plants for food and shelter.

TROPHIC STATE:

Trophic state is classification of lakes into categories of fertility and is a useful index rating and comparing lakes. From low to nutrient and algal biomass (as chlorophyll) concentrations, the trophic states are; **oligotrophic, mesotrophic, eutrophic** and **hypereutrophic (Table 2)**.

A majority of lakes in Alberta contain naturally high levels of chlorophyll *a* (8 to $\mu\text{g/L}$) due to our deep fertile soils. These are usually considered fertile and are termed eutrophic. The nutrient and algal biomass concentrations that define these categories shown in the following table, a figure of Alberta lakes compared by trophic state can found on the ALMS website.




Figure B: Suggested changes in various lake characteristics with eutrophication. From “Ecological Effects of Wastewater”, 1980.

Table A - Trophic status classification based on lake water characteristics.

Trophic state	Total Phosphorus ($\mu\text{g}\cdot\text{L}^{-1}$)	Total Nitrogen ($\mu\text{g}\cdot\text{L}^{-1}$)	Chlorophyll a ($\mu\text{g}\cdot\text{L}^{-1}$)	Secchi Depth (m)
Oligotrophic	< 10	< 350	< 3.5	> 4
Mesotrophic	10 – 30	350 - 650	3.5 - 9	4 - 2
Eutrophic	30 – 100	650 - 1200	9 - 25	2 - 1
Hypereutrophic	> 100	> 1200	> 25	< 1