



*The Alberta Lake Management Society  
Volunteer Lake Monitoring Program*

## Pigeon Lake Report

# 2018

Lakewatch is made possible  
with support from:





# ALBERTA LAKE MANAGEMENT SOCIETY'S LAKEWATCH PROGRAM

LakeWatch has several important objectives, one of which is to collect and interpret water quality data on Alberta Lakes. Equally important is educating lake users about their aquatic environment, encouraging public involvement in lake management, and facilitating cooperation and partnerships between government, industry, the scientific community and lake users. LakeWatch Reports are designed to summarize basic lake data in understandable terms for a lay audience and are not meant to be a complete synopsis of information about specific lakes. Additional information is available for many lakes that have been included in LakeWatch and readers requiring more information are encouraged to seek those sources.

ALMS would like to thank all who express interest in Alberta's aquatic environments and particularly those who have participated in the LakeWatch program. These leaders in stewardship give us hope that our water resources will not be the limiting factor in the health of our environment.

If you require the data in this report, please contact ALMS for the raw data files.



## ACKNOWLEDGEMENTS

The LakeWatch program is made possible through the dedication of its volunteers. A special thanks to Daren Lorentz for his commitment to collecting data at Pigeon Lake. We would also like to thank Alanna Robertson, Lindsay Boucher and Shona Derlukewich, who were summer technicians in 2018. Executive Director Bradley Peter and Program Coordinator Laura Redmond were instrumental in planning and organizing the field program. This report was prepared by Caitlin Mader and Bradley Peter.

## PIGEON LAKE

Pigeon Lake is a large (96.32 km<sup>2</sup>), shallow (average depth = 6m) lake located in the counties of Wetaskiwin and Leduc. It is a very popular recreational lake within easy driving distance from the cities of Edmonton, Leduc, and Wetaskiwin. Pigeon Lake lies within the Battle River watershed. Water flows into the lake through intermittent streams draining the west and northwest portions of the watershed. The outlet, Pigeon Lake Creek, at the southeast margin of the lake, drains toward the Battle River.<sup>1</sup> The lake's drainage basin is small (176.62 km<sup>2</sup>) but heavily developed with agriculture, oil and gas, and community developments throughout the watershed.<sup>2</sup>



Pigeon Lake at Poplar Bay 2017 (Photo by Laura Redmond)

The lake name is a translation from the Cree Mehmew Sâkâhikan, which means 'Dove Lake', but by 1858 the name Pigeon Lake was in use.<sup>3</sup> It has been suggested that the name Pigeon Lake refers to the huge flocks of Passenger Pigeons that once ranged in the area.<sup>1</sup> The lake was also previously known as Woodpecker Lake, and the Stoney name is recorded as Ke-gemni-wap-ta.<sup>3</sup> The water quality of Pigeon Lake is typical of large, productive, shallow lakes in Alberta, with water remaining quite green for most of the summer. However, residents have recently expressed concern over perceptions of deteriorating water quality as a result of recurring blue-green algal blooms, fish kills, and beach advisories<sup>4</sup>. Due to these concerns, there has been a demand to examine ways to reduce the frequency and intensity of cyanobacteria blooms. In 2013, data was collected to prepare a nutrient budget for Pigeon Lake - this report was later released in 2014 and it outlines areas of interest when considering watershed and in-lake management options<sup>5</sup>. In 2017 the Pigeon Lake Watershed Association released their draft Pigeon Lake Watershed Management Plan which can be accessed via [www.plwmp.ca](http://www.plwmp.ca).

The watershed area for Pigeon Lake is 176.62 km<sup>2</sup> and the lake area is 97.32 km<sup>2</sup>. The lake to watershed ratio of Pigeon Lake is 1:2. A map of the Pigeon Lake watershed area can be found <http://alms.ca/wp-content/uploads/2016/12/Pigeon.pdf>.

---

<sup>1</sup>Mitchell, P. and Prepas, E. (1990). Atlas of Alberta Lakes, University of Alberta Press. Retrieved from <http://sunsite.ualberta.ca/projects/alberta-lakes/>

<sup>2</sup> Teichreb, C., Peter, B. and Dyer, A. (2013). 2013 Overview of Pigeon lake Water Quality, Sediment Quality, and Non-Fish Biota. 2 pp.

<sup>3</sup> Aubrey, M. K. (2006). Concise place names of Alberta. Retrieved from <http://www.albertasource.ca/placenames/resources/searchcontent.php?book=1>

<sup>4</sup> Aquality Environmental Consulting. (2008). Pigeon Lake State of Watershed Report. Prepared for Pigeon Lake Watershed Alliance. Retrieved from: [www.plwa.ca](http://www.plwa.ca).

<sup>5</sup> Teichreb, C. (2014). Pigeon Lake Phosphorus Budget. Alberta Environment and Sustainable Resource Development. 28 pp.



## METHODS

**Profiles:** Profile data is measured at the deepest spot in the main basin of the lake. At the profile site, temperature, dissolved oxygen, pH, conductivity and redox potential are measured at 0.5- 1.0 m intervals. Additionally, Secchi depth is measured at the profile site and used to calculate the euphotic zone. On one visit per season, metals are collected at the profile site by hand grab from the surface and at some lakes, 1 m off bottom using a Kemmerer.

**Composite samples:** At 10-sites across the lake, water is collected from the euphotic zone and combined across sites into one composite sample. This water is collected for analysis of water chemistry, chlorophyll-a, nutrients and microcystin. Quality control (QC) data for total phosphorus was taken as a duplicate true split on one sampling date. ALMS uses the following accredited labs for analysis: Routine water chemistry and nutrients are analyzed by Maxxam Analytics, chlorophyll-a and metals are analyzed by Alberta Innotech, and microcystin is analyzed by the Alberta Centre for Toxicology (ACTF). In lakes where mercury samples are taken, they are analyzed by the Biogeochemical Analytical Service Laboratory (BASL).

**Invasive Species:** Monitoring for invasive quagga and zebra mussels involved two components: monitoring for juvenile mussel veligers using a 63 µm plankton net at three sample sites and monitoring for attached adult mussels using substrates installed at each lake.

**Data Storage and Analysis:** Data is stored in the Water Data System (WDS), a module of the Environmental Management System (EMS) run by Alberta Environment and Parks (AEP). Data goes through a complete validation process by ALMS and AEP. Users should use caution when comparing historical data, as sampling and laboratory techniques have changed over time (e.g. detection limits). For more information on data storage, see AEP Surface Water Quality Data Reports at [aep.alberta.ca/water](http://aep.alberta.ca/water).

Data analysis is done using the program R.<sup>1</sup> Data is reconfigured using packages tidyr<sup>2</sup> and dplyr<sup>3</sup> and figures are produced using the package ggplot2<sup>4</sup>. Trophic status for each lake is classified based on lake water characteristics using values from Nurnberg (1996)<sup>5</sup>. The Canadian Council for Ministers of the Environment (CCME) guidelines for the Protection of Aquatic Life are used to compare heavy metals and dissolved oxygen measurements. Pearson's Correlation tests are used to examine relationships between TP, chlorophyll-a, TKN and Secchi depth, providing a correlation coefficient (r) to show the strength (0-1) and a p-value to assess significance of the relationship.

---


<sup>1</sup> R Core Team (2016). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

<sup>2</sup> Wickman, H. and Henry, L. (2017). tidyr: Easily Tidy Data with 'spread ( )' and 'gather ( )' Functions. R package version 0.7.2. <https://CRAN.R-project.org/package=tidyr>.

<sup>3</sup> Wickman, H., Francois, R., Henry, L. and Muller, K. (2017). dplyr: A Grammar of Data Manipulation. R package version 0.7.4. <http://CRAN.R-project.org/package=dplyr>.

<sup>4</sup> Wickham, H. (2009). ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

<sup>5</sup> Nurnberg, G.K. (1996). Trophic state of clear and colored, soft- and hardwater lakes with special consideration of nutrients, anoxia, phytoplankton and fish. Lake and Reservoir Management 12: 432-447.



BEFORE READING THIS REPORT, CHECK  
OUT [A BRIEF INTRODUCTION TO  
LIMNOLOGY](#) AT [ALMS.CA/REPORTS](#)

## WATER CHEMISTRY

*ALMS measures a suite of water chemistry parameters. Phosphorus, nitrogen, and chlorophyll-*a* are important because they are indicators of eutrophication, or excess nutrients, which can lead to harmful algal/cyanobacteria blooms. One direct measure of harmful cyanobacteria blooms are Microcystins, a common group of toxins produced by cyanobacteria. See Table 2 for a complete list of parameters.*

The average total phosphorus (TP) concentration for Pigeon Lake was 32.8 µg/L (Table 2), falling just above threshold for eutrophic, or highly productive trophic classification. This value falls within the range of historical averages. Detected TP was lowest when first sampled in June at 19 µg/L, and peaked at 44 µg/L in late August (Figure 1).

Average chlorophyll-*a* concentrations in 2018 was 39.2 µg/L (Table 2), falling into the hypereutrophic, or very high productivity trophic classification. Chlorophyll-*a* was highest when first sampled in June at 73.5 µg/L, and reached its lowest point of 14.3 µg/L on July 19.

Finally, the average TKN concentration was 0.97 mg/L (Table 2) with concentrations peaking on August 31 at 1.2 mg/L.

Average pH was measured as 8.56 in 2018, buffered by moderate alkalinity (164 mg/L CaCO<sub>3</sub>) and bicarbonate (188 mg/L HCO<sub>3</sub>). Calcium was the dominant ion contributing to a low conductivity of 328 µS/cm (Table 2).




Figure 1- Total Phosphorus (TP), Total Kjeldahl Nitrogen (TKN), and Chlorophyll-*a* concentrations measured five times over the course of the summer at Pigeon Lake.

## METALS

*Samples were analyzed for metals once throughout the summer (Table 3). In total, 27 metals were sampled for. It should be noted that many metals are naturally present in aquatic environments due to the weathering of rocks and may only become toxic at higher levels.*

Metals were not measured in Pigeon Lake in 2018. Table 3 presents historical metal concentrations from previously measured years.

## WATER CLARITY AND SECCHI DEPTH

*Water clarity is influenced by suspended materials, both living and dead, as well as dissolved colored compounds in the water column. During the melting of snow and ice in spring, lake water can become turbid (cloudy) from silt transported into the lake. Lake water usually clears in late spring but then becomes more turbid with increased algal growth as the summer progresses. The easiest and most widely used measure of lake water clarity is the Secchi depth. Two times the Secchi depth equals the euphotic depth – the depth to which there is enough light for photosynthesis.*

The average Secchi depth of Pigeon Lake in 2018 was 2.29 m (Table 2). This depth was shallowest on August 10<sup>th</sup>, indicating low water clarity. Secchi depth was deepest on September 19<sup>th</sup>, indicating the highest water clarity.




Figure 2 – Secchi depth values measured five times over the course of the summer at Pigeon Lake in 2018.

## WATER TEMPERATURE AND DISSOLVED OXYGEN

*Water temperature and dissolved oxygen profiles in the water column can provide information on water quality and fish habitat. The depth of the thermocline is important in determining the depth to which dissolved oxygen from the surface can be mixed. Please refer to ALMS' [Brief Introduction to Limnology](#) for descriptions of technical terms.*

Temperatures of Pigeon Lake varied throughout the summer, with a minimum temperature of 9.8°C at 7m on September 19, and a maximum temperature of 21.4°C measured at the surface on August 10 (Figure 3a). The lake was not stratified during any of the sampling trips, with temperatures either constant or displaying a steady gradient from top to bottom. This indicates partial or complete mixing throughout the season.

Pigeon Lake remained well oxygenated through most of the water column throughout the summer, measuring above the CCME guidelines of 6.5 mg/L for the Protection of Aquatic Life (Figure 3b). The oxygen level fell below this level close to lake bottom on July 19 and August 10.




Figure 3 – a) Temperature (°C) and b) dissolved oxygen (mg/L) profiles for Pigeon Lake measured five times over the course of the summer of 2018.





## MICROCYSTIN

*Microcystins are toxins produced by cyanobacteria (blue-green algae) which, when ingested, can cause severe liver damage. Microcystins are produced by many species of cyanobacteria which are common to Alberta's Lakes, and are thought to be the one of the most common cyanobacteria toxins. In Alberta, recreational guidelines for microcystin are set at 20 µg/L. Blue-green algae advisories are managed by Alberta Health Services. Recreating in algal blooms, even if microcystin concentrations are not above guidelines, is not recommended.*

Composited microcystin levels in Pigeon Lake fell below the recreational guideline of 20 µg/L on each sampling date (Table 1). Caution should still be observed when recreating in visible cyanobacteria blooms.

Table 1 – Microcystin concentrations measured five times at Pigeon Lake in 2018.

Date	Microcystin Concentration (µg/L)
22-Jun-18	0.10
19-Jul-18	0.11
10-Aug-18	0.32
31-Aug-18	1.66
19-Sep-18	0.69
<b>Average</b>	0.58

## INVASIVE SPECIES MONITORING

*Dreissenid mussels pose a significant concern for Alberta because they impair the function of water conveyance infrastructure and adversely impact the aquatic environment. These invasive mussels have been linked to creating toxic algae blooms, decreasing the amount of nutrients needed for fish and other native species, and causing millions of dollars in annual costs for repair and maintenance of water-operated infrastructure and facilities.*

Monitoring involved two components: monitoring for juvenile mussels (veligers) using a plankton net and monitoring for attached adult mussels using substrates installed in each lake. No mussels have been detected in Pigeon Lake.

## WATER LEVELS

*There are many factors influencing water quantity. Some of these factors include the size of the lake's drainage basin, precipitation, evaporation, water consumption, ground water influences, and the efficiency of the outlet channel structure at removing water from the lake. Requests for water quantity monitoring should go through Alberta Environment and Parks Monitoring and Science division.*

Water levels in Pigeon Lake have remained relatively stable since Environment Canada began monitoring the lake in 1972 (Figure 4). Since 1972, Pigeon Lake water levels fluctuated between a maximum of 850.6 m asl and a minimum of 849.4 m asl. Data from Environment Canada was only available until 2015. A weir at the mouth of the outlet was installed in 1983 by ESRD to maintain water levels at 849.935 meters above sea level (m asl). In 2008, monitoring revealed that the weir had risen 0.15 m due to frost heaving. In 2013, the height was adjusted by ESRD to bring the structure back to the intended level of 849.935 m asl.<sup>1</sup>




Figure 4- Water levels measured in meters above sea level (m asl) from 1972-2015. Data retrieved from Environment Canada.

<sup>1</sup> Teichreb, C., Peter, B. and Dyer, A. (2013). 2013 Overview of Pigeon lake Water Quality, Sediment Quality, and Non-Fish Biota. 2 pp.

[illegible]

Table 2b: Average Secchi depth and water chemistry values for Pigeon Lake. Historical values are given for reference.

Parameter	1996	1997	1983	2003	2005	2006	2008	2010	2011	2013	2014
TP (µg/L)	37.5	29.9	35	63	26.5	60.3	26.3	40.7	74.7	27.3	45.8
TDP (µg/L)	/	/	/	/	6	38	9	13	19.1	7.69	16.4
Chlorophyll- <i>a</i> (µg/L)	18.5	12.77	15	36.9	9.2	21.9	7.98	21.92	66.2	12.28	19.24
Secchi depth (m)	1.8	2.5	1.5	1.38	1.9	2.7	4.42	2.75	1.25	3.23	2.31
TKN (mg/L)	/	/	0.611	1.075	0.71	1.1	0.67	1.033	1.5	0.78	0.72
NO <sub>2</sub> -N and NO <sub>3</sub> -N (µg/L)	/	/	1	/	3	29	13	7.67	15.9	5.91	26
NH <sub>3</sub> -N (µg/L)	/	/	3	/	2.5	124	16	72.3	108.9	28.39	24.6
DOC (mg/L)	/	/	/	/	/	7	/	7.35	/	/	8.3
Ca (mg/L)	/	/	/	/	28.85	21.13	27.2	23.75	19.5	27.62	22.83
Mg (mg/L)	/	/	/	/	12.65	14.12	12.87	13.85	12.5	12.84	11.43
Na (mg/L)	14.6	18.6	/	18.7	20	21	20.33	21.95	20.1	20.57	23.6
K (mg/L)	/	/	/	/	6.1	6.63	6.17	6.3	6.2	6.59	6.6
SO <sub>4</sub> <sup>2-</sup> (mg/L)	/	/	/	/	7.3	10.2	5.47	9	3.38	6.38	5.03
Cl <sup>-</sup> (mg/L)	/	/	/	/	4	3.33	3.33	3.05	3.03	3.19	3.5
CO <sub>3</sub> (mg/L)	/	/	/	/	8	4.67	3.33	0.5	8.7	3.27	5.92
HCO <sub>3</sub> (mg/L)	163	190	/	168.5	183	180	198	195	161	194.53	191.6
pH	8.66	8.17	/	8.56	8.6	8.5	8.37	8.57	8.74	8.34	8.59
Conductivity (µS/cm)	293	304	/	/	313	287	321.7	309.5	286.7	320	314
Hardness (mg/L)	106	130	/	103	125	119	121	116	100.2	122	104
TDS (mg/L)	151	169	/	/	176.5	173	175.33	173.5	153	176	182.3
Total Alkalinity (mg/L CaCO <sub>3</sub> )	149	156	/	151	163	155.3	165.7	160	146.7	164	157
Microcystin (µg/L)	/	/	/	/	/	/	/	0.087	0.173	0.1354	0.97

Table 2c: Average Secchi depth and water chemistry values for Pigeon Lake. Historical values are given for reference.

Parameter	2015	2016	2017	2018
TP ( $\mu\text{g/L}$ )	60.6	26	47.2	32.8
TDP ( $\mu\text{g/L}$ )	11.2	6	4.1	5.3
Chlorophyll- <i>a</i> ( $\mu\text{g/L}$ )	40.84	27.9	57.7	39.2
Secchi depth (m)	1.65	3.36	1.85	2.29
TKN (mg/L)	1.338	0.85	1.3	0.97
NO <sub>2</sub> -N and NO <sub>3</sub> -N ( $\mu\text{g/L}$ )	2.5	2.5	2.26	8.56
NH <sub>3</sub> -N ( $\mu\text{g/L}$ )	31	25	20.5	16.4
DOC (mg/L)	7.8	7.06	7.9	7.98
Ca (mg/L)	19.8	25.8	24.6	25.2
Mg (mg/L)	12.6	14.2	14.6	14.2
Na (mg/L)	20.8	23.8	24	23.6
K (mg/L)	6.1	7.4	7.34	7.34
SO <sub>4</sub> <sup>2-</sup> (mg/L)	4.5	6	5.36	6.94
Cl <sup>-</sup> (mg/L)	3.8	3.72	4.46	4.56
CO <sub>3</sub> (mg/L)	3.61	5.56	7.62	6.56
HCO <sub>3</sub> (mg/L)	178	184	178	188
pH	8.48	8.60	8.63	8.556
Conductivity ( $\mu\text{S/cm}$ )	298	320	316	328
Hardness (mg/L)	100.8	124	120	124
TDS (mg/L)	166	190	182	196
Total Alkalinity (mg/L CaCO <sub>3</sub> )	154	160	160	164
Microcystin ( $\mu\text{g/L}$ )	2.318	0.13	0.47	0.58

Table 3: Concentrations of metals were last measured in Pigeon Lake in August 2017. The CCME heavy metal Guidelines for the Protection of Freshwater Aquatic Life (unless otherwise indicated) are presented for reference.

Metals (Total Recoverable)	2003	2012	2014	2015	2016	2017	Guidelines
Aluminum µg/L	14.9	5.13	10.55	14.3	7.2	5.1	100 <sup>a</sup>
Antimony µg/L	0.05	0.06685	0.089	0.0785	0.066	0.064	/
Arsenic µg/L	1.67	1.375	2.285	2.145	2.06	1.87	5
Barium µg/L	78.5	89.75	77.35	74.1	73.5	78.1	/
Beryllium µg/L	0.02	0.00675	0.004	0.004	0.004	0.0015	100 <sup>c,d</sup>
Bismuth µg/L	0.0025	0.00125	0.0005	0.00325	0.001	0.0015	/
Boron µg/L	27.9	29.85	27.4	28.5	31.1	27.3	1500
Cadmium µg/L	0.01	0.00325	0.002	0.004	0.005	0.005	0.26 <sup>b</sup>
Chromium µg/L	0.27	0.015	0.5235	0.09	0.04	0.05	/
Cobalt µg/L	0.11	0.00605	0.006565	0.018	0.005	0.021	1000 <sup>d</sup>
Copper µg/L	1.08	0.2255	0.4155	0.235	0.5	0.22	4 <sup>b</sup>
Iron µg/L	39	2.04	15.75	144.2	20.6	14.3	300
Lead µg/L	0.145	0.0167	0.245	0.0595	0.028	0.032	7 <sup>b</sup>
Lithium µg/L	8.6	9.09	8.29	9.175	11.2	9.47	2500 <sup>e</sup>
Manganese µg/L	54.1	16.9	15.75	49.65	6.48	27.5	200 <sup>e</sup>
Molybdenum µg/L	0.62	0.704	0.731	0.728	0.907	0.711	73 <sup>c</sup>
Nickel µg/L	0.16	0.0025	0.3465	0.0205	0.219	0.88	150 <sup>b</sup>
Selenium µg/L	0.25	0.103	0.35	0.03	0.21	0.1	1
Silver µg/L	0.0025	0.0015	0.00681	0.002	0.003	0.0005	0.25
Strontium µg/L	245	234	261	233	249	231	/
Thallium µg/L	0.0015	0.00105	0.00291	0.000875	0.0075	0.003	0.8
Thorium µg/L	0.0015	0.008725	0.003575	0.011425	0.01	0.005	/
Tin µg/L	0.05	0.0549	0.0231	0.0355	0.027	0.03	/
Titanium µg/L	1.5	0.8925	1.4355	3.125	1	1.37	/
Uranium µg/L	0.086	0.1805	0.1945	0.167	0.161	0.144	15
Vanadium µg/L	0.26	0.1545	0.456	0.14	0.26	0.189	100 <sup>d,e</sup>
Zinc µg/L	1.5	0.899	1.56	0.65	0.9	0.3	30

Values represent means of total recoverable metal concentrations.

<sup>a</sup> Based on pH ≥ 6.5

<sup>b</sup> Based on water hardness > 180mg/L (as CaCO<sub>3</sub>)

<sup>c</sup> CCME interim value.

<sup>d</sup> Based on CCME Guidelines for Agricultural use (Livestock Watering).

<sup>e</sup> Based on CCME Guidelines for Agricultural Use (Irrigation).

A forward slash (/) indicates an absence of data or guidelines.



## LONG TERM TRENDS

Trend analysis was conducted on the parameters total phosphorus (TP), chlorophyll-*a*, total dissolved solids (TDS) and Secchi depth to look for changes over time in Pigeon Lake. In sum, non-significant increases were observed in TP and non-significant decreasing trends were observed in chlorophyll-*a* and Secchi depth. Secchi depth can be subjective and is sensitive to variation in weather - trend analysis must be interpreted with caution. TDS trend analysis was significant showing an increasing trend. Data is presented below as both a line graph (all data points) or a box-and-whisker plot. Detailed methods are available in the *ALMS Guide to Trend Analysis on Alberta Lakes*.

Table 1: Summary table of trend analysis on Pigeon Lake data from 2003 to 2018.

Parameter	Date Range	Trend	Probability
Total Phosphorus	1983-2018	Decreasing	Not significant
Chlorophyll- <i>a</i>	1983-2018	Increasing	Significant
Total Dissolved Solids	1983-2018	Increasing	Significant
Secchi Depth	1983-2018	Increasing	Significant

## Definitions:


**Median:** the value in a range of ordered numbers that falls in the middle.

**Trend:** a general direction in which something is changing.

**Monotonic trend:** a gradual change in a single direction.

**Statistically significant:** The likelihood that a relationship between variables is caused by something other than random chance. This is indicated by a *p*-value of <0.05. **Variability:** the extent by which data is inconsistent or scattered.

**Box and Whisker Plot:** a box-and-whisker plot, or boxplot, is a way of displaying all of our annual data. The median splits the data in half. The 75<sup>th</sup> percentile is the upper quartile of the data, and the 25<sup>th</sup> percentile is the lower quartile of the data. The top and bottom points are the largest and smallest observations.



## Total Phosphorus (TP)

Trend analysis of TP over time showed that it has not significantly changed in Pigeon Lake since 1982 (Tau = 0.09,  $p = 0.24$ ). Recent recent years show greater variation in total phosphorous levels than have been seen historically.




Figure 1- Monthly total phosphorus (TP) concentrations measured between June and September over the long term sampling dates between 1983 and 2018 (n = 90). The value closest to the 15<sup>th</sup> day of the month was chosen to represent the monthly value in cases with multiple monthly samples.

## Chlorophyll-*a*

Chlorophyll-*a* has increased significantly over time at Pigeon Lake ( $\text{Tau} = 0.22$ ,  $p = 0.052$ , slope =  $9.0 \times 10^{-4}$ ). The rate of change is very low, with the slope of the trend suggesting an increase of around  $3.29 \mu\text{g}/\text{decade}$  over the 35 years sampled. Chlorophyll-*a* trends follow TP trends with correlation over time ( $r = 0.66$ ,  $p = <0.001$ ).




Figure 2-Monthly chlorophyll-*a* concentrations measured between June and September over the long term sampling dates between 1983 and 2018 ( $n = 105$ ). The value closest to the 15<sup>th</sup> day of the month was chosen to represent the monthly value in cases with multiple monthly samples. Line graph is overlain by TP concentrations.

## Total Dissolved Solids (TDS)

Trend analysis showed a significantly increasing trend in TDS since 1983 in Pigeon Lake (Tau = 0.57,  $p = <0.001$ ).




Figure 3- Monthly TDS values measured between June and September over the long term sampling dates between 1983 and 2018 ( $n = 91$ ). The value closest to the 15<sup>th</sup> day of the month was chosen to represent the monthly value in cases with multiple monthly samples.

## Secchi Depth

Secchi depth has been increasing significantly in Pigeon Lake since 1983, but at a very slow rate of change ( $\text{Tau} = 0.16$ ,  $p = 0.01$ ). Recent years show much greater variation both within and between seasons.




Figure 4- Monthly Secchi depth values measured between June and September over the long term sampling dates between 1983 and 2018 ( $n = 69$ ). The value closest to the 15<sup>th</sup> day of the month was chosen to represent the monthly value in cases with multiple monthly samples.

Table 2- Results of trend tests using total phosphorus (TP), chlorophyll-*a*, total dissolved solids (TDS) and Secchi depth data from June to September on Pigeon Lake data.

Definition	Unit	Total Phosphorus (TP)	Chlorophyll-a	Total Dissolved Solids (TDS)	Secchi Depth
Statistical Method	-	Mann Kendall	Mann Kendall	Seasonal Kendall	Seasonal Kendall
The strength and direction (+ or -) of the trend between -1 and 1	Tau	0.085	0.19	0.57	0.16
The extent of the trend	Slope	0.0004	0.0009	0.90	0.01
The statistic used to find significance of the trend	Z	1.19	2.86	6.30	2.04
Number of samples included	n	90	105	91	69
The significance of the trend	<i>p</i>	0.24	<0.004*	<0.001*	0.04*

\**p* < 0.05 is significant within 95%